

RANDOM POLYNOMIALS WITH PRESCRIBED NEWTON POLYTOPE, I

BERNARD SHIFFMAN AND STEVE ZELDITCH

ABSTRACT. The Newton polytope P_f of a polynomial f is well known to have a strong impact on its zeros, as in the Kouchnirenko-Bernstein theorem on the number of simultaneous zeros of m polynomials with given Newton polytopes. In this article, we show that P_f also has a strong impact on the distribution of zeros of one or several polynomials. We equip the space of (holomorphic) polynomials of degree $\leq N$ in m complex variables with its usual $SU(m+1)$ -invariant Gaussian measure and then consider the conditional measures $\gamma_{|NP}$ induced on the subspace of polynomials whose Newton polytope $P_f \subset NP$. When $P = \Sigma$, the unit simplex, then it is obvious and well-known that the expected distribution of zeros $Z_{f_1, \dots, f_k}$ (regarded as a current) of k polynomials $f_1, \dots, f_k$ of degree N is uniform relative to the Fubini-Study form. Our main results concern the conditional expectation $\mathbf{E}_{|NP}(Z_{f_1, \dots, f_k})$ of zeros of k polynomials with Newton polytope $NP \subset Np\Sigma$ (where $p = \deg P$); these results are asymptotic as the scaling factor $N \rightarrow \infty$. We show that $\mathbf{E}_{|NP}(Z_{f_1, \dots, f_k})$ is asymptotically uniform on the inverse image $\mathcal{A}_P$ of the open scaled polytope $\frac{1}{p}P^\circ$ via the moment map $\mu : \mathbb{CP}^m \rightarrow \Sigma$ for projective space. However, the zeros have an exotic distribution outside of $\mathcal{A}_P$ and when $k = m$ (the case of the Kouchnirenko-Bernstein theorem) the percentage of zeros outside $\mathcal{A}_P$ approaches 0 as $N \rightarrow \infty$.

CONTENTS

Introduction	2
1. Background on toric varieties and moment polytopes	10
1.1. Newton polytopes and toric varieties	10
1.2. Szegő kernels	12
1.3. Moment maps and torus actions	15
2. Szegő kernels on toric varieties	17
2.1. The exact formula	17
2.2. Toeplitz operator approach to Π_N^{MP}	19
3. Mass asymptotics	24
3.1. The classically allowed region	27
3.2. The classically forbidden region	28
4. Distribution of Zeros	42
4.1. Expected distribution of zeros and Szegő kernels	42
4.2. Asymptotics of zeros	45
4.3. Amoebas in the plane	48
5. Examples	50
5.1. The square	50
5.2. The Hirzebruch surfaces	51
5.3. Remarks on the behavior at transition points	53
References	55

Date: March 12, 2002.

Research partially supported by NSF grant DMS-0100474 (first author) and DMS-0071358 (second author).

INTRODUCTION

It is well known that the Newton polytope P_f of a holomorphic polynomial $f(z_1, \dots, z_m)$ of degree p has a crucial influence on its value distribution and in particular on its zero set. Even the number of simultaneous zeros in $\mathbb{C}^{*m} := (\mathbb{C} \setminus \{0\})^m$ of m generic polynomials $f_1, \dots, f_m$ depends on their Newton polytopes P_{f_j} [Be, Ko1, Ko2]. Our purpose in this paper is to demonstrate that the Newton polytope of a polynomial f also has a crucial influence on its *mass density* $|f(z)|^2 dV$ and on the *spatial distribution* of zeros $\{f = 0\}$. We will show that there is a *classically allowed region* where the mass almost surely concentrates and a *classically forbidden region* where it almost surely is exponentially decaying. The classically allowed region is the inverse image $\mu_\Sigma^{-1}(\frac{1}{p}P)$ of the (scaled) polytope $\frac{1}{p}P$ under the standard moment map μ_Σ of $\mathbb{CP}^m$. The simultaneous zeros of m generic polynomials $f_1, \dots, f_m$ in $\mathbb{C}^{*m}$ almost surely concentrates (in the limit of high degrees) in the classically allowed region, giving a kind of quantitative version of the Bernstein-Kouchnirenko theorem. The Newton polytope has an equally strong (though different) impact for $k < m$ polynomials $f_1, \dots, f_k$. The image of the zero set of $f_1, \dots, f_k$ under the moment map is (up to a logarithmic re-parametrization) known as an *amoeba* in the sense of [GKZ, Mi2]. Results on the expected distribution of amoebas can be obtained from our results on the expected zero current; for example, as a consequence of Corollary 6, for polytopes $\frac{1}{p}P$ with vertices in the interior of the standard simplex $\Sigma \subset \mathbb{R}^m$, there is also a *very forbidden region* which the amoeba almost surely avoids. These patterns in zeros are statistical—they hold for random polynomials with prescribed Newton polytope—and are asymptotic as the degree of the polynomials tends to infinity.

To state our problems and results precisely, let us recall some definitions. Let

$$f(z_1, \dots, z_m) = \sum_{\alpha \in \mathbb{N}^m: |\alpha| \leq p} c_\alpha \chi_\alpha(z_1, \dots, z_m), \quad \chi_\alpha(z) = z_1^{\alpha_1} \cdots z_m^{\alpha_m}$$

be a polynomial of degree p in m complex variables. By the *support* of f we mean the set

$$S_f = \{\alpha \in \mathbb{Z}^m : c_\alpha \neq 0\}, \quad (1)$$

and by its *Newton polytope* P_f we mean the integral polytope

$$P_f := \hat{S}_f = \text{the convex hull in } \mathbb{R}^m \text{ of } S_f. \quad (2)$$

Throughout this paper we will make the simplifying assumption that P is a Delzant polytope [De, Gu1]. Non-Delzant polytopes are equally important in the study of polynomials, and we plan to extend our results to more general convex integral polytopes in the sequel [SZ2]. We also plan to study a parallel tunneling phenomenon for critical points.

Our aim is to study the statistical patterns in polynomials with a fixed Newton polytope. We therefore define a natural Gaussian measure $d\gamma_P$ on the space of polynomials with support contained in a fixed Newton polytope P . Since our purpose is to compare zero sets and masses as the polytope P varies, we view the polytope as placing a condition on the Gaussian ensemble of all polynomials of degree p equal to that of P , i.e. we define the measures $d\gamma_P$ as conditional probability measures of one fixed Gaussian measure. To define this ensemble, we homogenize the polynomial f to obtain a homogeneous polynomial F of degree p in $m+1$ complex variables. Such polynomials may be identified with holomorphic sections $F \in H^0(\mathbb{CP}^m, \mathcal{O}(p))$ of the p^{th} power of the hyperplane section bundle $\mathcal{O}_{\mathbb{CP}^m}(1)$. We denote by

$$H^0(\mathbb{CP}^m, \mathcal{O}(p), P) = \{F \in H^0(\mathbb{CP}^m, \mathcal{O}(p)) : P_f \subset P\} \quad (3)$$

the space of homogeneous polynomials F of degree d whose associated inhomogeneous form $f(z_1, \dots, z_m) = F(1, z_1, \dots, z_m)$ has Newton polytope P_f contained in P .

Let us recall that $H^0(\mathbb{CP}^m, \mathcal{O}(p))$ carries the standard $\text{SU}(m+1)$ -invariant inner product

$$\langle F_1, \bar{F}_2 \rangle = \int_{S^{2m+1}} F_1 \bar{F}_2 d\sigma,$$

where $d\sigma$ is Haar measure on the $(2m+1)$ -sphere S^{2m+1} . (See (28) for an alternate description in terms of the Fubini-Study form on $\mathbb{CP}^m$ and the Fubini-Study metric on $\mathcal{O}(1)$.) An orthonormal basis of $H^0(\mathbb{CP}^m, \mathcal{O}(p))$

is given by $\{\|\chi_\alpha\|^{-1}\chi_\alpha\}_{|\alpha|\leq p}$, where $\|\cdot\|$ denotes the norm in $H^0(\mathbb{CP}^m, \mathcal{O}(p))$. (Note that $\|\chi_\alpha\|$ depends on p .) The corresponding $\mathrm{SU}(m+1)$ -invariant Gaussian measure γ_p is defined by

$$d\gamma_p(s) = \frac{1}{\pi^{k_p}} e^{-|s|^2} ds, \quad s = \sum_{|\alpha|\leq p} \lambda_\alpha \frac{\chi_\alpha}{\|\chi_\alpha\|}, \quad (4)$$

where $k_p = \#\{\alpha : |\alpha| \leq p\} = \binom{m+p}{p}$. Thus, the coefficients λ_α are independent complex Gaussian random variables with mean zero and variance one. We then endow the space $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ with the associated conditional probability measure $\gamma_p|_P$:

$$d\gamma_p|_P(s) = \frac{1}{\pi^{\#P}} e^{-|s|^2} ds, \quad s = \sum_{\alpha \in P} \lambda_\alpha \frac{\chi_\alpha}{\|\chi_\alpha\|}, \quad (5)$$

where the coefficients λ_α are again independent complex Gaussian random variables with mean zero and variance one. (By a slight abuse of notation, we let $\sum_{\alpha \in P}$ denote the sum over the lattice points $\alpha \in P \cap \mathbb{Z}^m$; $\#P$ denotes the cardinality of $P \cap \mathbb{Z}^m$.) We observe that, as a subspace of $H^0(\mathbb{CP}^m, \mathcal{O}(p))$, $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ inherits the inner product $\langle s_1, s_2 \rangle$ and that $\gamma|_P$ is the induced Gaussian measure. Probabilities relative to $\gamma|_P$ can be considered as conditional probabilities; i.e. for any event E ,

$$\mathrm{Prob}_\gamma\{f \in E | P_f = P\} = \mathrm{Prob}_{\gamma|_P}(E).$$

We now wish to consider the expected distribution of mass and zeros of polynomials with fixed Newton polytope. It turns out to involve the moment map μ_Σ of projective space. We use a notation for this moment map which reflects its dependence on the standard unit simplex Σ in $\mathbb{R}^m$ with vertices at $(0, \dots, 0)$, $(1, 0, \dots, 0)$, $(0, 1, \dots, 0)$, $\dots$, $(0, \dots, 0, 1)$. The ‘projective moment map’ is then given by

$$\mu_\Sigma(z) = \left(\frac{|z_1|^2}{1 + \|z\|^2}, \dots, \frac{|z_m|^2}{1 + \|z\|^2} \right).$$

Via this moment map we define:

Definition: Let $P \subset \mathbb{R}_+^m$ be an integral polytope. The classically allowed region for polynomials in $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ is the set

$$\mathcal{A}_P := \mu_\Sigma^{-1} \left(\frac{1}{p} P^\circ \right) \subset \mathbb{C}^{*m}$$

(where P° denotes the interior of P), and the classically forbidden region is its complement $\mathbb{C}^{*m} \setminus \mathcal{A}_P$.

Our first result concerns the simultaneous zero set of m independent polynomials in m variables. Bézout’s theorem tells us that m generic homogeneous polynomials $F_1, \dots, F_m$ of degree p have exactly p^m simultaneous zeros; these zeros all lie in $\mathbb{C}^{*m}$, for generic F_j . In fact, one immediately sees (by uniqueness of Haar probability measure) that the expected distribution of zeros is uniform with respect to the Fubini-Study projective volume form, when the ensemble is given the $\mathrm{SU}(m+1)$ -invariant measure $d\gamma_p$.

According to Kouchnirenko’s theorem [Ko1, Ko2, At, Au], the number of joint zeros in $\mathbb{C}^{*m}$ of m generic polynomials $\{f_1, \dots, f_m\}$ with given Newton polytope P equals $m! \mathrm{Vol}(P)$, where $\mathrm{Vol}(P)$ is the Euclidean volume of P . For example, if $P = p\Sigma$, where Σ is the standard unit simplex in $\mathbb{R}^m$, then $\mathrm{Vol}(p\Sigma) = p^m \mathrm{Vol}(\Sigma) = \frac{p^m}{m!}$, and we get Bézout’s theorem. (More generally, the f_j may have different Newton polytopes, in which case the number of zeros is given by the Bernstein-Kouchnirenko formula as a ‘mixed volume.’)

Now consider m independent random polynomials with Newton polytope P , using the conditional probability (5). We let $\mathbf{E}_{|P}(Z_{f_1, \dots, f_m})$ denote the expected density of their simultaneous zeros. It is the measure on $\mathbb{C}^{*m}$ given by

$$\mathbf{E}_{|P}(Z_{f_1, \dots, f_m})(U) = \int d\gamma_p|_P(f_1) \cdots \int d\gamma_p|_P(f_m) [\#\{z \in U : f_1(z) = \cdots = f_m(z) = 0\}], \quad (6)$$

for $U \subset \mathbb{C}^{*m}$, where the integrals are over $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$. In fact, $\mathbf{E}_{|P}(Z_{f_1, \dots, f_m})$ is an absolutely continuous measure given by a $\mathcal{C}^\infty$ density (see Prop. 4.2). Our first result is the surprising fact that as the polytope P expands, these zeros are expected to concentrate in the classically allowed region and are (asymptotically) uniform there:

THEOREM 1. *Suppose that P is a Delzant polytope in $\mathbb{R}^m$. Then*

$$\frac{1}{(Np)^m} \mathbf{E}_{|NP}(Z_{f_1, \dots, f_m}) \rightarrow \begin{cases} \omega_{\text{FS}}^m & \text{on } \mathcal{A}_P \\ 0 & \text{on } \mathbb{C}^{*m} \setminus \mathcal{A}_P \end{cases},$$

*in the distribution sense; i.e., for any open $U \subset \mathbb{C}^{*m}$, we have*

$$\frac{1}{(Np)^m} \mathbf{E}_{|NP}(\#\{z \in U : f_1(z) = \dots = f_m(z) = 0\}) \rightarrow m! \text{Vol}_{\mathbb{CP}^m}(U \cap \mathcal{A}_P).$$

In fact, our methods imply that the convergence of the zero current on the classically allowed region is exponentially fast in the sense that

$$\mathbf{E}_{|NP}(Z_{f_1, \dots, f_m}) = (Np)^m \omega_{\text{FS}}^m + O(e^{-\lambda N}) \quad \text{on } \mathcal{A}_P,$$

for some positive continuous function λ on $\mathcal{A}_P$.

Theorem 1 is a special case of a general result stated below (Theorem 4) on the expected distribution of simultaneous zeros of k random polynomials with given Newton polytopes $P_1, \dots, P_k$. Although we only discuss expected behavior of zeros here, it is possible to show (using methods similar to those of [SZ1]) that distribution of zeros is self-averaging: i.e., almost all polynomials exhibit the expected behavior in an asymptotic sense. For the sake of brevity, we concentrate here on only the essentially new aspects of our problem, namely the dependence of the expected behavior on P .

The following illustration¹ shows the classically allowed region (shaded) and the classically forbidden region (unshaded) when P is the unit square in $\mathbb{R}^2$ (and $p = 2$):




FIGURE 1. The classically allowed region for $P = [0, 1] \times [0, 1]$

The terminology is taken from the semiclassical analysis of ground states of Schroedinger operators $H_h = -\hbar^2 \Delta + V$ on $\mathbb{R}^n$. The well-known ‘Agmon estimates’ of ground states (cf. [Ag]) show that $\mathcal{L}^2$ -normalized ground states or low-lying eigenfunctions $H_h \varphi = E \varphi$ of H are concentrated as $\hbar \rightarrow 0$ in the classically allowed region $C_E = \{x \in \mathbb{R}^n : V(x) \leq E\}$ and have exponential decay $|\varphi(x)|^2 = O(e^{-d(x, C_E)/\hbar})$ as $\hbar \rightarrow 0$ in the complement. Here, $d(x, C_E)$ is the ‘Agmon distance’ to the classically allowed region, i.e. the distance in the metric $\sqrt{V(x) - E} dx$. In our setting, the Hamiltonian is $\bar{\partial}^* \bar{\partial}$ on $\mathcal{L}^2$ -sections of powers $\mathcal{O}(Np)$ of the hyperplane section bundle, $\hbar = 1/N$, and the ground states are the holomorphic sections $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$.

¹We would like to thank Aaron Carass for providing the illustrations in Figures 1, 3, 4, 6 and 7.

Replacing $|\varphi(x)|^2$ is the expected ‘mass density’ of the polynomials in this subspace. By definition, the mass density of a polynomial f at a point $z \in \mathbb{C}^{*m}$ is the Fubini-Study norm

$$|f(z)|_{\text{FS}} := \frac{|f(z)|}{(1 + \|z\|^2)^{\delta/2}} \quad (\delta = \deg f). \quad (7)$$

Our next result describes the asymptotics of the expected mass density of $\mathcal{L}^2$ -normalized polynomials with Newton polytope NP , i.e. the expected density $\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2)$ with respect to the Haar probability measure, denoted by ν_{NP} , on the unit sphere in $H^0(\mathbb{CP}^m, \mathcal{O}(NP), NP)$. One easily shows (see §3) that

$$\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2) = \frac{1}{\#(NP)} \mathbf{E}_{|NP}(|f(z)|_{\text{FS}}^2) = \frac{1}{\#(NP)} \Pi_{|NP}(z, z), \quad (8)$$

where $\Pi_{|NP}$ is the Szegő kernel of $H^0(\mathbb{CP}^m, \mathcal{O}(NP), NP)$. Here, the term ‘Szegő kernel’ of a space $\mathcal{S}$ of holomorphic sections of a line bundle refers to the kernel for the orthogonal projection to $\mathcal{S}$ from the space $\mathcal{L}^2(\partial D)$ of square-integrable functions on the boundary of the associated unit disk bundle D ; i.e., it is of the form $\Pi(x, y) = \sum_j s_j(x) \overline{s_j(y)}$, where $\{s_j\}$ is an orthonormal basis of $\mathcal{S}$. Precise definitions and the relationship to the classical Szegő kernel for a strictly pseudoconvex domain are described in §1.2. Note that

$$\#(NP) = \dim H^0(\mathbb{CP}^m, \mathcal{O}(NP), NP) = \text{Vol}(P)N^m + \dots,$$

which is a polynomial given by the Riemann-Roch formula (see §3).

We shall show that the expected mass density $\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2)$ is asymptotically uniform with respect to Fubini-Study measure in the classically allowed region (as if there were no constraint at all); while in the forbidden region the mass decays exponentially. Thus in the semiclassical limit $N \rightarrow \infty$, all the mass concentrates in the classically allowed region. To describe the behavior in the forbidden region, we subdivide it into regions corresponding to the faces of the polytope $\frac{1}{p}P$ that lie in the interior Σ° of the simplex. To describe these regions, we recall that the fan associated to $\frac{1}{p}P$ (or P) is the collection of polyhedral cones $C_F \subset \mathbb{R}^m$ normal to the faces F of $\frac{1}{p}P$ (see §1.1.1). For each face $F \subset \Sigma^\circ$, we let

$$\mathcal{R}_F = \{e^\tau \cdot z : \tau \in C_F, \mu_\Sigma(z) \in F\}. \quad (9)$$

Note that if F is the open face $\frac{1}{p}P^\circ$, then $\mathcal{R}_F$ is the classically allowed region $\mathcal{A}_P$. We show in §3.2.1 that the classically forbidden region decomposes into the (disjoint) forbidden subregions

$$\left\{ \mathcal{R}_F : F \text{ is a face of } \frac{1}{p}P, F \subset \Sigma^\circ, F \neq \frac{1}{p}P^\circ \right\}.$$

We call $\mathcal{R}_F$ the *flow-out* of the face F . Boundary points of the $\mathcal{R}_F$ are called *transition points*. Points of $\partial\mathcal{A}_P$ are always transition points, but there may be others.

For the case where P is the unit square, then there are two interior faces whose flow-outs are the connected components of the classically forbidden region (see Figure 1 above). In this case, the set of transition points equals $\partial\mathcal{A}_P$. We shall give an example with an interior vertex, where the forbidden region is connected but decomposes into flow-outs of 3 faces (see Figure 7 in §5.2.1); in this case there are transition points not in $\partial\mathcal{A}_P$.

THEOREM 2. *Suppose that P is a Delzant polytope in $\mathbb{R}^m$. Then the expected mass density of random $\mathcal{L}^2$ normalized polynomials with Newton polytope NP has $\mathcal{C}^\infty$ asymptotic expansions of the form:*

$$\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2) \sim \begin{cases} \frac{\prod_{j=1}^m (pN+j)}{\#(NP)} = c_0 + c_1 N^{-1} + c_2 N^{-2} + \dots, & \text{for } z \in \mathcal{A}_P = \mu_\Sigma^{-1}(\frac{1}{p}P^\circ) \\ N^{-s/2} e^{-Nb(z)} [c_0^F(z) + c_1^F(z)N^{-1} + c_2^F(z)N^{-2} + \dots], & \text{for } z \in \mathcal{R}_F^\circ \end{cases}$$

where c_0^F and $b|_{\mathcal{R}_F^\circ}$ are positive $\mathcal{C}^\infty$ functions on $\mathcal{R}_F^\circ$, and $s = \text{codim } F$ (for each face $F \subset \Sigma^\circ$). Furthermore, the remainder estimates in the expansions are uniform on compact subsets of the open regions $\mathcal{R}_F^\circ$ and of $\mathcal{A}_P$.

We shall use (8) to restate Theorem 2 in terms of asymptotic formulae for the conditional Szegő kernel on the diagonal for which we give more precise asymptotics in Proposition 3.1. In [SZ2] we will extend the result to the non-Delzant case as well. Heuristically, the mass concentration in $\mathcal{A}_P$ can be understood as follows: the mass of a monomial m_α concentrates (exponentially) on the torus $\mu_\Sigma^{-1}(\alpha)$. The constraint $\alpha \in NP$ thus

concentrates all the mass of polynomials with fixed Newton polytope NP in $\mathcal{A}_P$, with exponentially small errors, and the mass there is uniformly distributed with small errors.

In particular,

$$\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2) \longrightarrow \begin{cases} \frac{p^m}{\text{Vol}(P)}, & \text{for } z \in \mathcal{A}_P \\ 0, & \text{for } z \in \mathbb{C}^{*m} \setminus \overline{\mathcal{A}_P} \end{cases},$$

as illustrated by the following graphs, plotted using Maple, for the case where P is the unit square. (Recall Figure 1 for the depiction of $\mathcal{A}_P$ for this case.)




FIGURE 2. $\frac{1}{4}\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2)$ for $P = [0, 1] \times [0, 1]$

The decay function b in Theorem 2 is analogous to the Agmon action to the allowed region. We shall show in Proposition 3.1 that b is $\mathcal{C}^1$ (but not $\mathcal{C}^2$) on all of $\mathbb{C}^{*m}$, and we derive several formulas for $b(z)$ ((10), (11), (116)). To describe our first formula, we associate with each point $z \in \mathbb{C}^{*m} \setminus \mathcal{A}_P$ a unique point $\xi \in \partial\mathcal{A}_P$ of the form $\xi = e^{\tau/2} \cdot z$, where $-\tau$ is in the (real) normal cone to the convex set P at the point $p\mu_\Sigma(\xi) \in \partial P$. Here $r \cdot z = (r_1 z_1, \dots, r_m z_m)$ denotes the $\mathbb{R}_+^m$ action on $\mathbb{C}^{*m}$. (The normal cone to a point $q \in \partial P$, described in §1.1.1, is the element of the fan associated to the face of P containing q .) We write $\tau_z = \tau$, $q(z) = p\mu_\Sigma(\xi)$; these vectors are then given by the conditions:

- $p\mu_\Sigma(e^{\tau_z/2} \cdot z) = q(z) \in \partial P$;
- $-\tau_z$ is in the normal cone to P at $q(z)$.

The existence and uniqueness of $q(z)$ and τ_z are stated in Lemma 3.8.

Our first formula for b is :

$$b(z) = \langle q(z), \tau_z \rangle + p \log \left(\frac{1 + \|z\|^2}{1 + \|e^{\tau_z/2} \cdot z\|^2} \right) \quad \text{for } z \in \mathbb{C}^{*m} \setminus \mathcal{A}_P. \quad (10)$$

We will obtain (10) from the integral formula,

$$b(z) = \int_0^{\tau_z} \left[q(e^{\sigma/2} \cdot z) - p\mu_\Sigma(e^{\sigma/2} \cdot z) \right] \cdot d\sigma \quad (11)$$

(where $\int_0^{\tau_z}$ denotes the integral over any path in $\mathbb{R}^m$ from 0 to τ_z), which could be interpreted as an action, thereby bringing the results closer to classical Agmon estimates.

We also show that $\frac{1}{N} \log \mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2) \rightarrow -b(z)$ uniformly on compact subsets of $\mathbb{C}^{*m}$ (Proposition 3.2).

Our final results concern the zero set of one or more random polynomials. We shall consider not only the expected volume density of the zero set, but also the more precise description involving the expected zero current. The zero current $Z_{f_1, \dots, f_k}$ of k polynomials (or more generally, k holomorphic functions) $f_1, \dots, f_k$ is the current of integration over the zero set

$$|Z_{f_1, \dots, f_k}| := \{z \in \mathbb{C}^{*m} : f_1(z) = \dots = f_m(z) = 0\}.$$

This current is given by

$$(Z_{f_1, \dots, f_k}, \varphi) := \int_{|Z_{f_1, \dots, f_k}|} \varphi, \quad \text{for a test form } \varphi \in \mathcal{D}^{m-k, m-k}(\mathbb{C}^{*m}),$$

where integration is over the set of regular points of $|Z_{f_1, \dots, f_k}|$ (provided that it is of codimension k and without multiplicity, which is almost surely the case).

We denote by $\mathbf{E}_{|P}(Z_f) = \mathbf{E}_{\gamma_P|P}(Z_f)$ the conditional expectation of the zero current of a random polynomial $f \in H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ with Newton polytope P . In fact, $\mathbf{E}_{|P}(Z_f)$ is actually a smooth $(1, 1)$ -form on $\mathbb{C}^{*m}$ (Proposition 4.1).

Let us recall what happens when $P = p\Sigma$ (cf. [SZ1]). By the uniqueness of Haar measure, the expected zero current $\mathbf{E}(Z_f)$ taken over all polynomials of degree p is given by $p\omega_{\text{FS}}$, where $\omega_{\text{FS}} = \frac{i}{2\pi} \partial \bar{\partial} \log \|z\|^2$ is the Fubini-Study Kähler form on $\mathbb{CP}^m$. Thus the expected distribution of zeros, as well as the tangent to the zero varieties, is uniform over $\mathbb{CP}^m$. We now consider how the expectation changes if we add the condition that $P_f = P$.

THEOREM 3. *Let P be a Delzant polytope. Then there exists a closed semipositive $(1, 1)$ -form ψ_P on $\mathbb{C}^{*m}$ with piecewise $\mathcal{C}^\infty$ coefficients such that:*

- i) $N^{-1}\mathbf{E}_{|NP}(Z_f) \rightarrow \psi_P$ in $\mathcal{L}_{\text{loc}}^1(\mathbb{C}^{*m})$.
- ii) $\psi_P = p\omega_{\text{FS}}$ on the classically allowed region $\mu_\Sigma^{-1}(\frac{1}{p}P^\circ)$.
- iii) On each region $\mathcal{R}_F^\circ$, the $(1, 1)$ -form ψ_P is $\mathcal{C}^\infty$ and has constant rank equal to $\dim F$; in particular, if $v \in \Sigma^\circ$ is a vertex of $\frac{1}{p}P$, then $\psi_P|_{\mathcal{R}_v^\circ} = 0$.

As a corollary, we obtain some statistical results on the so-called ‘tentacles’ of amoebas in dimension 2 (see §4.3). Roughly speaking, the (compact) amoeba of a polynomial $f(z_1, z_2)$ is the image of the Riemann surface Z_f under the moment map μ_Σ on $(\mathbb{C}^*)^2$, and the tentacles are the ends of the amoeba. Certain tentacles must end at vertices of the triangle Σ while others are ‘free’ to end anywhere along the boundary of Σ . In Corollary 4.7, we will prove that (in the limit $N \rightarrow \infty$) almost all of the free tentacles of typical amoebas tend to end in the classically allowed portion of $\partial\Sigma$.

We call the form ψ_P in Theorem 3 the *limit expected zero current*. By $\mathcal{L}_{\text{loc}}^1$ convergence in (i), we mean $\mathcal{L}_{\text{loc}}^1$ convergence of the coefficients. (Recall that $\mathbf{E}_{|NP}(Z_{f_1, \dots, f_k})$ is a (k, k) -form with smooth coefficients.) If we write $\psi_P = \sqrt{-1} \sum \psi_{jk}(z) dz_j \wedge d\bar{z}_k$, then $(\psi_{jk}(z))$ is a semi-positive Hermitian matrix, for non-transition points z . By the rank of ψ_P at z , we mean the rank of the matrix $(\psi_{jk}(z))$. Note that if $\mathcal{R}_F$ and $\mathcal{R}_{F'}$ are adjoining regions (i.e., have a common codimension 1 interface), then F and F' are of different dimensions, so ψ_P must be discontinuous along the interface. We remark that as an element of $\mathcal{D}'^{1,1}(\mathbb{C}^{*m})$, the current ψ_P is closed and positive.

The form ψ_P not only encodes the expected (normalized) density of the zero set, but also the expected density of tangent directions to the zero set. In the course of the proof of Theorem 3, we will show that in the forbidden region, the limit tangent directions are restricted. In particular, as the polytope expands, the tangent spaces to typical zero sets through a point $z^0 \in \mathcal{R}_F^\circ$ approach spaces containing the tangent space (at z^0) of the ‘normal flow’ $\{e^{\tau+i\theta} \cdot z^0 : \tau, \theta \in T_F^\perp \subset \mathbb{R}^m\}$. Precise formulations of this fact are given in Theorem 4.5 and (162). Thus, while the expected distribution of zero densities is absolutely continuous, the expected distribution of zero tangents is singular.

We can also consider k independently chosen random polynomials

$$f_j \in H^0(\mathbb{CP}^m, \mathcal{O}(p_j), P_j), \quad 1 \leq j \leq k$$

($k \leq m$), and we let $\mathbf{E}_{|P_1, \dots, P_k}(Z_{f_1, \dots, f_k})$ denote the expected zero current with respect to the probability measure $\gamma_{p_1|P_1} \times \dots \times \gamma_{p_k|P_k}$ on the product space. If $P_1 = \dots = P_k = P$, then we also write $E_{|P, \dots, P}(Z_{f_1, \dots, f_k}) = E_{|P}(Z_{f_1, \dots, f_k})$.

THEOREM 4. *Let $P_1, \dots, P_k$ be Delzant polytopes. Then*

$$N^{-k} \mathbf{E}_{|NP_1, \dots, NP_k}(Z_{f_1, \dots, f_k}) \rightarrow \psi_{P_1} \wedge \dots \wedge \psi_{P_k} \quad \text{in } \mathcal{L}_{\text{loc}}^1(\mathbb{C}^{*m}), \quad \text{as } N \rightarrow \infty.$$

We see from Theorem 3(iii) that the zero set $|Z_f|$ of a polynomial with polytope NP almost surely creeps into the classically forbidden region $\mu_\Sigma^{-1}(\Sigma \setminus \frac{1}{p}P)$ in the semiclassical limit $N \rightarrow \infty$. Indeed the expected volume of the zero set, or more generally the simultaneous zero set of k polynomials, has the following exotic distribution law:

COROLLARY 5. *Let $P_1, \dots, P_k$ be Delzant polytopes. Then for any open set $U \subset \subset \mathbb{C}^{*m}$,*

$$\frac{1}{N^k} \mathbf{E}_{|NP_1, \dots, NP_k} \text{Vol}(|Z_{f_1, \dots, f_k}| \cap U) \rightarrow \frac{1}{(m-k)!} \int_U \psi_{P_1} \wedge \dots \wedge \psi_{P_k} \wedge \omega_{\text{FS}}^{m-k}.$$

By the volume of $|Z_{f_1, \dots, f_k}| \cap U$, we mean the $(2m-2k)$ -dimensional volume in $\mathbb{CP}^m$, which is given by

$$\text{Vol}(|Z_{f_1, \dots, f_k}| \cap U) = \int_{|Z_{f_1, \dots, f_k}| \cap U} \frac{1}{(m-k)!} \omega^{m-k} = \left(Z_{f_1, \dots, f_k}, \frac{1}{(m-k)!} \omega^{m-k} \right). \quad (12)$$

Corollary 5 follows immediately from Theorem 4 and (12).

In addition, the following consequence of Corollary 5 and part (iii) of Theorem 3 says that there are subtler ‘very forbidden regions’ in the case where the polytope has faces of codimension ≥ 2 in the interior of Σ , namely the regions comprising the normal flow-out of these faces:

COROLLARY 6. *The expected zero current $N^{-k} \mathbf{E}_{|NP}(Z_{f_1, \dots, f_k})$ tends to 0 at all points of each forbidden subregion $\mathcal{R}_F$ with $\dim F < k$. In particular, $N^{-1} \mathbf{E}_{|NP}(Z_f)$ tends to zero at all points of the flow-out $\mathcal{R}_v$ of a vertex $v \in \frac{1}{p}P \cap \Sigma^\circ$.*

We now say a few words about the proofs. The key result is Theorem 2 on the mass of polynomials with Newton polytope P , which we formulate more precisely in Proposition 3.1 using the conditional Szegő kernel. As will be shown in §4, we can derive the expected distribution of zeros from our asymptotic formula for the conditional Szegő kernel $\Pi_{|PN}(z, z)$.

As in the proof in [At, Au] of Kouchnirenko’s theorem, a natural approach to our results on polynomials with prescribed Newton polytope is through the Kähler toric variety (M_P, ω_P) associated to P . We recall that M_P carries a natural line bundle L_P with $c_1(L_P) = \omega_P$. The main connection is that there exists a natural identification

$$H^0(\mathbb{CP}^m, \mathcal{O}(Np), P) \simeq H^0(M_P, L_P^N) \quad (13)$$

between polynomials with prescribed Newton polytope and holomorphic sections of powers L_P^N of L_P . The torus $\mathbf{T}^m = \{(e^{i\varphi_1}, \dots, e^{i\varphi_m})\}$ acts on $H^0(M_P, L_P^N)$ with character

$$\chi_{NP}(e^{i\varphi}) = \sum_{\alpha \in NP} e^{i\langle \alpha, \varphi \rangle}, \quad e^{i\varphi} = (e^{i\varphi_1}, \dots, e^{i\varphi_m}). \quad (14)$$

We then have the simple expression

$$\Pi_{|NP}(z, z) = \int_{\mathbf{T}^m} \Pi_{NP}^{\mathbb{CP}^m}(t \cdot z, z) \overline{\chi_{NP}(t)} dt. \quad (15)$$

To obtain asymptotics as $N \rightarrow \infty$, we derive an oscillatory integral formula. First we combine (15) with the formula

$$\chi_{NP}(e^{i\varphi}) = \int_{M_P} \Pi_{NP}^{M_P}(e^{i\varphi} \cdot w, w) d\text{Vol}_{M_P}(w), \quad (16)$$

where $\Pi_{NP}^{M_P}$ is the Szegő kernel of M_P , i.e. the orthogonal projection onto $H^0(M_P, L_P^N)$ with respect to the volume form determined by ω_P . In order to obtain an exponential decay rate $e^{-Nb(z)}$ of Theorem 2, we will

need a very precise understanding of $\Pi_N^{M_P}$. This we obtain in §2 by making a special construction of the Szegő kernel of a toric variety which is perhaps of independent interest. Our formula for Π^{M_P} involves the application of a certain *Toeplitz Fourier multiplier* to the classical Szegő kernel for the sphere, pulled back by a monomial embedding. After showing in Proposition 2.2 that this Toeplitz-Fourier multiplier is a Toeplitz operator in the sense of [BG], we obtain an oscillatory integral formula for $\Pi_{|NP}$ (see (69) and (72)) in terms of an integral over $\mathbf{T}^m \times M_P$ with complex phase of positive type.

After that, the proof of Theorem 2 follows from the method of stationary phase for complex oscillatory integrals. For the case where z is in the classically allowed region, we easily find that the critical manifold is given by $\varphi = 0$ and $\mu_P(w) = p\mu_\Sigma(z)$, where μ_P is the moment map of M_P . Since the phase vanishes and has nondegenerate normal Hessian along the critical manifold, we immediately obtain an asymptotic expansion. The case where z is in the classically forbidden region is more subtle. Since $p\mu_\Sigma(z)$ lies outside of P for this case, the phase has no critical points. To complete the analysis, we must deform the contour to pick up critical points. The main part of the proof of Theorem 2 involves finding a contour for which there is a nondegenerate critical manifold where the phase has maximum real part. In particular, we consider the complexification $\mathbb{C}^{*m}$ of $\mathbf{T}^m$ and deform $\mathbf{T}^m$ to a contour of the form $(\log |\zeta_1|, \dots, \log |\zeta_m|) = \tau \in \mathbb{R}^m$. We then show that the phase takes its maximal real part on a critical manifold if and only if τ is the unique vector τ_z used in formula (10) for the decay rate $b(z)$; indeed the maximal real part of the phase is $-b(z)$. Finally we prove that nondegeneracy of the normal Hessian holds if (and only if) z is not a transition point, so that we obtain the asymptotic expansion of Theorem 2.

We would like to remark here that other formulae for the lattice sums $\chi_P(t)$ are well known from the work of Khovanskii-Pukhlikov [KP], Brion-Vergne [BV] and others. An alternative approach is to replace (16) with the formula from [BV, Theorem 3.12]:

$$\chi_P(e^{i\varphi}) = \text{Todd}(\partial/\partial h) \left(\int_{P(h)} e^{i\langle x, \varphi \rangle} dx \right) \Big|_{h=0}, \quad (17)$$

where $P(h) = \{x : \langle u_j, x \rangle + a_j + h_j \geq 0, 1 \leq j \leq n\}$, and $\text{Todd}(\partial/\partial h)$ is a certain infinite order differential operator known as a *Todd operator*. Upon dilating the polytope, one obtains

$$\chi_{NP}(e^{i\varphi}) = N^m \text{Todd}(N\partial/\partial h) \left(\int_{P(h)} e^{iN\langle x, \varphi \rangle} dx \right) \Big|_{h=0}. \quad (18)$$

The asymptotics of mass and zeros in the classically allowed region follow very easily from this expression for χ_{NP} , and with further effort one can prove the result in the forbidden region. Both approaches seem to us quite interesting and to have their own advantages. In the sequel [SZ2], we will use the approach based on (18), which seems more efficient when the toric varieties have singularities.

We end the introduction with some comments on the relation of our results to other work on polynomials with a fixed Newton polytope. Such polynomials are called *sparse* in the literature, and methods of algebraic (including toric) geometry have recently been applied to the computational problem of locating zeros of systems of such sparse polynomials (e.g., see [HS, MR, St, Ro, Ve]). Our results give information on the expected location of zeros when the polynomials are given the conditional measure $\gamma_{|NP}$ on $H^0(\mathbb{CP}^m, \mathcal{O}(N), P)$. To our knowledge, the asymptotic concentration of zeros of sparse systems in the classically allowed region has not been observed before. It obviously pays most to search for zeros of systems of m polynomials in m variables in the allowed region.

It should be noted that this asymptotic pattern reflects the choice of the conditional measures $\gamma_{|NP}$ on $H^0(\mathbb{CP}^m, \mathcal{O}(N), P)$. In previous work [SZ1], we obtained the expected and almost everywhere distribution of zeros of several holomorphic sections of positive line bundles $L \rightarrow M$ over general Kähler manifolds. Those results specialize to our current setting where $M = M_P, L = L_P$ and also give results on the distribution of zeros of polynomials in $H^0(\mathbb{CP}^m, \mathcal{O}(N), P)$. However, in that paper we defined different Gaussian probability measures $\gamma_N^{M_P}$ on $H^0(M_P, L_P^N)$, namely those induced by the $\mathcal{L}^2$ inner product induced by a hermitian metric on L_P and its curvature form ω_P . In Proposition 4.6 of [SZ1], we showed that the expected distribution of zeros relative to the $\gamma_N^{M_P}$ satisfies $\frac{1}{N^k} \mathbf{E}_{\gamma_N^{M_P}}(Z_{f_1, \dots, f_k}) = \omega_P^k + O(\frac{1}{N})$. The measures $\gamma_N^{M_P}$ have more recently been studied by [MR], who also obtain (among other things) the formula for the expected distribution. Our

present measures $\gamma_N^{M_P}$ are quite singular relative to $\gamma_{|NP}$ in the limit as $N \rightarrow \infty$. Our point of view is that the conditional measures $\gamma_{|NP}$ on $H^0(\mathbb{CP}^m, \mathcal{O}(N), P)$ are the natural ones when comparing polynomials in $H^0(\mathbb{CP}^m, \mathcal{O}(N), P)$ as P varies. The toric variety M_P is a technical device for studying $\Pi_{|NP}$ but does not figure into the statements of our main results. The difference in the two measures is that in the conditional case, we fix the $\mathcal{L}^2$ norms of the monomials z^α in advance (as their $\mathbb{CP}^m$ norms), while in the $\gamma_N^{M_P}$ case the norms vary as P and M_P vary. The latter would seem to create complicated biases towards some monomials and away from others as P varies and make it difficult to understand what such a comparison is measuring.

1. BACKGROUND ON TORIC VARIETIES AND MOMENT POLYTOPES

1.1. Newton polytopes and toric varieties. The space $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ of polynomials with Newton polytope P may be identified with the space $H^0(M_P, L_P)$ of holomorphic sections of a line bundle $L_P \rightarrow M_P$ over a toric variety M_P . Under our running assumption that P is Delzant (see below), the variety M_P is smooth. Recall that a *toric variety* is a complex algebraic variety M containing the complex torus

$$\mathbb{C}^{*m} = (\mathbb{C} \setminus \{0\}) \times \cdots \times (\mathbb{C} \setminus \{0\})$$

as a Zariski-dense open set such that the group action of $\mathbb{C}^{*m}$ on itself extends to a $\mathbb{C}^{*m}$ action on M . In the Delzant case, M_P can be given the structure of a symplectic manifold such that the restriction of the action to the underlying real torus

$$\mathbf{T}^m = \{(\zeta_1, \dots, \zeta_m) \in \mathbb{C}^{*m} : |\zeta_j| = 1, 1 \leq j \leq m\}$$

is a Hamiltonian action (see §1.3).

We now review two (well-known) constructions of M_P , by fans, resp. monomial embeddings, to establish notation and clarify the properties we will be needing. For further details, see [Da, Fu, Oda]. We point out here that there is also a geometric approach due to Delzant [De] through *symplectic reduction*; see [De, Au, Gu1].

1.1.1. Fans. By a *convex integral polytope*, we mean the convex hull in $\mathbb{R}^n$ of a finite set in the lattice $\mathbb{Z}^m$. A convex integral polytope P with n *facets* (i.e. codimension-one faces) can be defined by linear equations

$$\ell_i(x) := \langle x, u_i \rangle + a_i \geq 0, \quad (i = 1, \dots, n),$$

where $u_i \in \mathbb{Z}^m$ is the primitive interior normal to the i -th facet. The polytope P is called *Delzant* if each vertex is the intersection of exactly m facets whose primitive normal vectors generate the lattice $\mathbb{Z}^m$.

For each point $x \in P$, we consider the *normal cone to P at x* ,

$$C_x := \{u \in \mathbb{R}^m : \langle u, x \rangle = \sup_{y \in P} \langle u, y \rangle\},$$

which is a closed convex polyhedral cone. We decompose P into a finite union of *faces*, each face being an equivalence class under the equivalence relation $x \sim y \iff C_x = C_y$. For each face F , we let C_F denote the normal cone of the points of F . **Note that by our convention, the faces are disjoint sets.** We shall use the term *closed face* to refer to the closure of a face of P .

Each face of dimension r ($0 \leq r \leq m$) is an open polytope in an r -plane in $\mathbb{R}^m$; i.e., the 0-dimensional faces are the vertices of P , the 1-dimensional faces are the edges with their end points removed, and so forth. The facets F_i and their normal cones are given by:

$$\bar{F}_i = \{x \in P : \ell_i(x) = 0\}, \quad C_{F_i} = \{-tu_i : t \geq 0\}.$$

The m -dimensional face is the interior P° of the polytope with normal cone $C_{P^\circ} = \{0\}$.

A convex integral polytope P determines the fan $\mathcal{F}_P := \{C_F : F \text{ is a face of } P\}$. A *fan* $\mathcal{F}$ in $\mathbb{R}^m$ is a collection of closed convex rational polyhedral cones such that a closed face of a cone in $\mathcal{F}$ is an element of $\mathcal{F}$ and the intersection of two cones in $\mathcal{F}$ is a closed face of each of them. An example of an integral polytope and its fan is given in the illustration below.

There is a natural way to construct a toric variety from a fan by gluing together the affine varieties arising from the cones in the fan. As we will not use this construction here, we refer the reader to [Fu] for the details. We note that a fan whose union is all of $\mathbb{R}^m$ defines a complete toric variety, which is projective if and only if $\mathcal{F}$ is the fan of a convex polytope P . If P is Delzant, then the toric variety of the fan $\mathcal{F}_P$ is a




FIGURE 3. A convex polytope and its fan

smooth projective manifold. The polytope in Figure 3 is Delzant if $w_3 = (0, 1)$, and the corresponding toric variety M_P is a Hirzebruch surface (see [Fu]). We shall study this example in §5.2.1.

1.1.2. *Monomial embeddings.* An alternative definition of M_P is through $\mathbf{T}^m$ -equivariant monomial embeddings. We let $P \cap \mathbb{Z} = \{\alpha(1), \alpha(2), \dots, \alpha(\#P)\}$. If we fix $c = (c_1, \dots, c_{\#P}) \in (\mathbb{C}^*)^{\#P}$, we get a map

$$\Phi_P^c = [c_{\alpha(1)}\chi_{\alpha(1)}, \dots, c_{\alpha(\#P)}\chi_{\alpha(\#P)}] : \mathbb{C}^{*m} \rightarrow \mathbb{CP}^{\#P-1} ;$$

i.e.,

$$\Phi_P^c(z) = [c_{\alpha(1)}z^{\alpha(1)}, \dots, c_{\alpha(\#P)}z^{\alpha(\#P)}] .$$

The closure of the image is a toric variety $M_P^c \subset \mathbb{CP}^{\#P-1}$. If P is Delzant, then Φ_P^c is injective and we identify $\mathbb{C}^{*m}$ with its image (the ‘open orbit’) in M_P^c .

We refer to the resulting embedding

$$\Phi_P^c : M_P^c \hookrightarrow \mathbb{CP}^{\#P-1} \quad (19)$$

as a monomial embedding, and we define the line bundle

$$L_P := \Phi_P^{c*} \mathcal{O}(1). \quad (20)$$

Furthermore, $H^0(M_P^c, L_P) = \Phi_P^{c*} H^0(\mathbb{CP}^{\#P-1}, \mathcal{O}(1))$; i.e., the sections are spanned by the monomials χ_α . It follows that

$$H^0(M_P^c, L_P) \simeq H^0(\mathbb{CP}^m, \mathcal{O}(p), P) . \quad (21)$$

We further have (see [Fu]):

$$H^0(M_P^c, L_P^N) \simeq \Phi_P^{c*} H^0(\mathbb{CP}^{\#P-1}, \mathcal{O}(N)) = H^0(\mathbb{CP}^m, \mathcal{O}(Np), NP) . \quad (22)$$

Recall that $H^0(\mathbb{CP}^{\#P-1}, \mathcal{O}(1))$ has as a basis the linear coordinate functions $\lambda_j : \mathbb{C}^{\#P} \rightarrow \mathbb{C}$, $1 \leq j \leq \#P$. The *Fubini-study metric* h_{FS} on $\mathcal{O}(1)$ is given by

$$|\lambda_j|_{\text{FS}}([\zeta]) = \frac{|\zeta_j|}{\|\zeta\|} \quad (\zeta \in \mathbb{C}^{\#P}) ,$$

which has curvature form $\omega_{\text{FS}} = \frac{i}{2\pi} \partial \bar{\partial} \log \|\zeta\|^2$. We endow L_P^c with the Hermitian metric $h_P^c := \Phi_P^{c*} h_{\text{FS}}$ of curvature ω_P^c given on $\mathbb{C}^{*m}$ by

$$\omega_P^c = \Phi_P^{c*} \omega_{\text{FS}} = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \sum_{\alpha \in P} |c_\alpha|^2 |z^\alpha|^2. \quad (23)$$

Each monomial χ_α with $\alpha \in P$ corresponds to a section of $H^0(M_P^c, L_P^c)$ and vice versa. To explicitly define this correspondence, we make the identifications (recalling (20)):

$$\chi_{\alpha(j)} \equiv c_{\alpha(j)}^{-1} \Phi_P^c * \zeta_j \in H^0(M_P^c, L_P^c) , \quad 1 \leq j \leq \#P . \quad (24)$$

So far, we have not specified the constants c_α . For studying our phenomena, the choice of constants defining the toric variety M_P is not important. However, when our polytope P is the full simplex $p\Sigma$, we shall use the special choice $c_\alpha = \binom{p}{\alpha}^{1/2}$, where $\binom{p}{\alpha}$ is the multinomial coefficient defined in the next section. (We use $c_\alpha = 1$ in §2, but except for Proposition 2.1, which is not used in our proof, any choice of c will work.)

Remark: For a fixed polytope P , the symplectic manifolds (M_P^c, ω_P^c) are equivariantly symplectically equivalent, by Delzant's theorem [De, Gu1, Au]. The M_P^c are also equivariantly holomorphically equivalent. However, in general the maps defining the equivalences are not simultaneously holomorphic and symplectic (see [Gu1]).

1.2. Szegő kernels. We first describe a general Szegő kernel for a positive Hermitian line bundle (L, h) on a compact complex manifold M . Let $\mathcal{S}_N \subset H^0(M, L^N)$ denote a subspace of the (finite dimensional) vector space of holomorphic sections of the N^{th} power of L . We give $\mathcal{S}_N$ an inner product. An important example is the ‘induced inner product’:

$$\langle s_1, \bar{s}_2 \rangle_{\mathcal{S}_N} = \int_M \left\langle s_1(z), \overline{s_2(z)} \right\rangle_{h_N} d\text{Vol}_M(z), \quad s_1, s_2 \in \mathcal{S}_N. \quad (25)$$

By the ‘Szegő projector’ $\Pi_{\mathcal{S}_N}$, we mean the orthogonal projection onto $\mathcal{S}_N$. As in [Ze, SZ1, BSZ], we lift our Szegő kernels on the associated principal S^1 bundle $X \rightarrow M$, defined via a Hermitian metric h as follows: Let $L^* \rightarrow M$ denote the dual line bundle to L with dual metric h^* , and put $X = \{v \in L^* : \|v\|_{h^*} = 1\}$. We let $r_\theta x = e^{i\theta} x$ ($x \in X$) denote the S^1 action on X . We then identify sections s_N of L^N with equivariant functions $\hat{s}_N$ on X by the rule

$$\hat{s}_N(\lambda) = (\lambda^{\otimes N}, s_N(z)), \quad \lambda \in X_z, \quad (26)$$

where $\lambda^{\otimes N} = \lambda \otimes \cdots \otimes \lambda$; then $\hat{s}_N(r_\theta x) = e^{iN\theta} \hat{s}_N(x)$, and we regard elements of $\mathcal{S}_N$ as equivariant functions on X .

If $\{S_j^N\}$ denotes an orthonormal basis of $\mathcal{S}_N$, then the projector $\Pi_{\mathcal{S}_N}$ is given by the kernel

$$\Pi_{\mathcal{S}_N}(x, y) = \sum_{j=1}^{k_N} \hat{S}_j^N(z) \overline{\hat{S}_j^N(y)}. \quad (27)$$

(Note that the usual Szegő kernel for the strictly pseudoconvex boundary X is given by $\sum_{N=1}^{\infty} \Pi_{\mathcal{S}_N}$, where $\mathcal{S}_N = H^0(M, L^N)$.)

We now describe three different sequences of Szegő kernels on toric varieties, which play a crucial role in our main results:

1.2.1. The projective Szegő kernels. As a first example, we let L be the hyperplane section bundle $\mathcal{O}(1)$ on $\mathbb{CP}^m$. Recall that the space $H^0(\mathbb{CP}^m, \mathcal{O}(p))$ of holomorphic sections of $L^p = \mathcal{O}(p)$ consists of homogeneous polynomials

$$F(\zeta_0, \dots, \zeta_m) = \sum_{|\lambda|=p} C_\lambda \zeta^\lambda \quad (\zeta^\lambda = \zeta_0^{\lambda_0} \cdots \zeta_m^{\lambda_m})$$

in $m+1$ variables. We can identify F with the (non-homogeneous) polynomial in m variables,

$$f(z_1, \dots, z_m) = F(1, z_1, \dots, z_m) = \sum_{|\alpha| \leq p} c_\alpha z^\alpha \quad (z^\alpha = z_1^{\alpha_1} \cdots z_m^{\alpha_m}),$$

where $c_\alpha = C_{\hat{\alpha}^p}$, $\hat{\alpha}^p = (p - |\alpha|, \alpha_1, \dots, \alpha_m)$. Inversely, a polynomial $f(z)$ of degree $\leq p$ can be identified with an element of $H^0(\mathbb{CP}^m, \mathcal{O}(p))$.

We give $\mathbb{CP}^m$ the Fubini-Study Kähler form given in homogeneous coordinates $(\zeta_0, \dots, \zeta_m)$ by $\omega_{\text{FS}} = \frac{i}{2\pi} \partial \bar{\partial} \log \|\zeta\|^2$, and we give $\mathcal{O}(p)$ the *Fubini-Study metric*:

$$|F(\zeta)|_{\text{FS}} = |F(\zeta)| / \|\zeta\|^p, \quad \text{for } F \in H^0(\mathbb{CP}^m, \mathcal{O}(p)).$$

Identifying F with the polynomial $f(z) = F(1, z_1, \dots, z_m)$, the Fubini-Study norm can be written

$$|f(z)|_{\text{FS}} = |f(z)| / (1 + \|z\|^2)^{p/2} \quad (z \in \mathbb{C}^m).$$

We equip the space $\mathcal{S}_p = H^0(\mathbb{CP}^m, \mathcal{O}(p))$ of all homogeneous polynomials of degree p with the inner product:

$$\langle f, \bar{g} \rangle = \int_{\mathbb{CP}^m} \langle F, \bar{G} \rangle_{\text{FS}} d\text{Vol}_{\mathbb{CP}^m} = \frac{1}{m!} \int_{\mathbb{C}^m} \frac{\langle f(z), \overline{g(z)} \rangle}{(1 + \|z\|^2)^p} \omega_{\text{FS}}^m(z), \quad f, g \in H^0(\mathbb{CP}^m, \mathcal{O}(p)). \quad (28)$$

(We use here the Riemannian volume $d\text{Vol}_{\mathbb{CP}^m} = \frac{1}{m!} \omega_{\text{FS}}^m$; note that the total volume of $\mathbb{CP}^m$ is $\frac{1}{m!}$, using our conventions.)

Under the $\mathbf{T}^m$ action, we have the weight space decomposition

$$H^0(\mathbb{CP}^m, \mathcal{O}(p)) = \bigoplus_{|\alpha| \leq p} \mathbb{C} \chi_\alpha,$$

where we recall that $\chi_\alpha(z) = z_1^{\alpha_1} \cdots z_m^{\alpha_m}$. The monomials $\{\chi_\alpha\}$ are orthogonal but not normalized. Any choice of norming constants $\{r_\alpha \in \mathbb{C}\}$ will give a monomial basis $\{r_\alpha \chi_\alpha\}$ for $H^0(\mathbb{CP}^m, \mathcal{O}(p))$. We shall choose $r_\alpha = \frac{1}{\|\chi_\alpha\|}$, where

$$\|\chi_\alpha\| = \sqrt{\langle \chi_\alpha, \chi_\alpha \rangle} = \left[\frac{p!}{(p+m)! \binom{p}{\alpha}} \right]^{\frac{1}{2}}, \quad \binom{p}{\alpha} := \frac{p!}{(p-|\alpha|)! \alpha_1! \cdots \alpha_m!},$$

is the Fubini-Study $\mathcal{L}^2$ norm of χ_α given by (28). (See [SZ1, §4.2]; the extra factor $m!$ in [SZ1] is due to the use of ω^m instead of $\frac{\omega^m}{m!}$ for the volume form.) This choice provides an orthonormal basis for $H^0(\mathbb{CP}^m, \mathcal{O}(p))$ given by the monomials

$$\frac{1}{\|\chi_\alpha\|} \chi_\alpha = \sqrt{\frac{(p+m)!}{p!} \binom{p}{\alpha}} \chi_\alpha, \quad |\alpha| \leq p.$$

We can identify $L^* = \mathcal{O}_{\mathbb{CP}^m}(-1)$ with $\mathbb{C}^{m+1}$ with the origin blown up, and the circle bundle $X \subset L^*$ is unit sphere $S^{2m+1} \subset \mathbb{C}^{m+1}$. We let $\widehat{\chi}_\alpha : S^{2m+1} \rightarrow \mathbb{C}$ denote the equivariant lift of $\chi_\alpha \in H^0(\mathbb{CP}^m, \mathcal{O}(p))$. We note that

$$\widehat{\chi}_\alpha(x) = x^{\hat{\alpha}^p},$$

and hence the Szegő kernel $\Pi_p^{\mathbb{CP}^m}$ for the orthogonal projection is given by:

$$\Pi_p^{\mathbb{CP}^m}(x, y) = \sum_{|\alpha| \leq p} \frac{1}{\|\chi_\alpha\|} \widehat{\chi}_\alpha(x) \overline{\widehat{\chi}_\alpha(y)} = \frac{(p+m)!}{p!} \sum_{|\alpha| \leq p} \binom{p}{\alpha} x^{\hat{\alpha}^p} \bar{y}^{\hat{\beta}^p} = \frac{(p+m)!}{p!} \langle x, \bar{y} \rangle^p, \quad (29)$$

for $x, y \in S^{2m+1}$. (The sum $\sum_{p=0}^\infty \Pi_p^{\mathbb{CP}^m}$ is the usual Szegő kernel for the sphere.)

1.2.2. The conditional Szegő kernels associated to a polytope P . In this case, the relevant space of polynomials is the subspace $H^0(\mathbb{CP}^m, \mathcal{O}(p), P) \subset H^0(\mathbb{CP}^m, \mathcal{O}(p))$ of polynomials with Newton polytope P . Here, we may choose any $p \geq \deg P := \max\{|\alpha| : \alpha \in P\}$, but we normally choose $p = \deg P$. In the conditional Szegő kernel, we retain the Fubini-Study inner product, denoted $\langle \cdot, \cdot \rangle_P$, on this subspace. Hence this example is very similar to the previous one. The main difference is that under the $\mathbf{T}^m$ action, we have the weight space decomposition

$$H^0(\mathbb{CP}^m, \mathcal{O}(p), P) = \bigoplus_{\alpha \in P} \mathbb{C} \chi_\alpha.$$

Definition: The conditional Szegő kernel Π_P is the kernel for the orthogonal projection to $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ with respect to the induced Fubini-Study inner product:

$$\Pi_P(x, y) = \sum_{\alpha \in P} \frac{1}{\|\chi_\alpha\|^2} \widehat{\chi}_\alpha(x) \overline{\widehat{\chi}_\alpha(y)}. \quad (30)$$

When defining the term ‘random polynomial with fixed Newton polytope P ’, we wish to use an $\mathcal{L}^2$ -norm on monomials which is defined independently of P . This explains why the conditional Szegő kernel is the

essential one in our problem. The conditional Szegő kernel can be written explicitly as

$$\Pi|_P(z, \theta; w, \varphi) = \frac{e^{ip(\theta-\varphi)} \sum_{\alpha \in P} \binom{p}{\alpha} z^\alpha \bar{w}^\alpha}{(1 + \|z\|^2)^{p/2} (1 + \|w\|^2)^{p/2}}, \quad (31)$$

where (z, θ) , resp. (w, φ) , are local coordinates for x , resp. y , in X ; i.e.,

$$x = e^{i\theta} (1 + \|z\|^2)^{-1/2} (1, z_1, \dots, z_m).$$

1.2.3. *The intrinsic Szegő kernels of a toric variety.* We now consider Szegő kernels which are defined by intrinsic inner products associated to the toric variety $M_P = M_P^c$. (To simplify notation, we now drop the c .) We regard them as a technical device for obtaining properties of the conditional Szegő kernels.

We thus put $\mathcal{S}_N = H^0(M_P, L_P^N)$, equipped with the inner product $\langle \cdot, \cdot \rangle_{M_P}$ derived from a Hermitian inner product h_P on L_P . As on all S^1 bundles associated to line bundles, the S^1 action on X_P gives the decompositions

$$\mathcal{L}^2(X_P) = \bigoplus_{N=1}^{\infty} \mathcal{L}_N^2(X_P), \quad f \in \mathcal{L}_N^2 \iff f(e^{i\theta} \cdot x) = e^{iN\theta} f(x).$$

Restricting to the Hardy space, we obtain

$$\mathcal{H}^2(X_P) = \bigoplus_{N=1}^{\infty} \mathcal{H}_N^2(X_P), \quad \mathcal{H}_N^2 = \mathcal{L}_N^2 \cap \mathcal{H}^2.$$

Recall that we have the canonical identifications $\mathcal{H}_N^2(X_P) \equiv H^0(M_P, L_P^N)$ given by identifying a section s_N with the equivariant function $\hat{s}_N$ on X_P . We let $\langle \cdot, \cdot \rangle_{X_P}$ denote the corresponding inner product on $\mathcal{H}_N^2(X_P)$, and we consider the orthogonal projectors $\Pi_N^{M_P}(x, y)$ relative to this inner product.

We let $\hat{\chi}_\alpha^P$ denote the equivariant lift of χ_α to X_P . Since $\langle \cdot, \cdot \rangle_{X_P}$ is invariant under the $\mathbf{T}^m$ action on X_P , the $\{\hat{\chi}_\alpha^P\}$ are orthogonal: indeed,

$$\langle \hat{\chi}_\alpha^P, \hat{\chi}_\beta^P \rangle_{X_P} = \langle t^* \hat{\chi}_\alpha^P, t^* \hat{\chi}_\beta^P \rangle_{X_P} = t^\alpha \bar{t}^\beta \langle \hat{\chi}_\alpha^P, \hat{\chi}_\beta^P \rangle_{X_P}, \quad \text{for all } t \in \mathbf{T}^m,$$

and hence $\langle \hat{\chi}_\alpha^P, \hat{\chi}_\beta^P \rangle_{X_P} = 0$ unless $\alpha = \beta$. We then have:

$$\Pi_N^{M_P}(x, y) = \sum_{\alpha \in NP} \frac{1}{\|\hat{\chi}_\alpha\|_{X_P}^2} \hat{\chi}_\alpha^P(x) \overline{\hat{\chi}_\alpha^P(y)}, \quad (32)$$

where $\|\hat{\chi}_\alpha\|_{X_P}$ is the $\mathcal{L}^2$ norm of $\hat{\chi}_\alpha$ with respect to the inner product $\langle \cdot, \cdot \rangle_{X_P}$.

In particular, we note that $M_\Sigma = M_{p\Sigma} = \mathbb{CP}^m$, and we have the circle bundles $X_{p\Sigma} \rightarrow \mathbb{CP}^m$, for $p \geq 1$. When $p = 1$, $X_\Sigma = S^{2m+1}$ while for $p > 1$, it is the lens space $X_{p\Sigma} = S^{2m+1}/\{e^{2\pi i/p}\}$. The latter statement follows from the fact that homogeneous polynomials of degree p are well defined on (and separate points of) the quotient by the cyclic group of p^{th} roots of unity.

1.2.4. *Powers of a line bundle.* Our main results are asymptotics results in powers of a line bundle or (equivalently) dilates of the polytope P . We therefore take powers of the line bundles in the previous three examples obtaining the spaces $H^0(\mathbb{CP}^m, \mathcal{O}(Np), NP)$ and $H^0(M_P, L_P^N) \simeq H^0(M_{NP}, L_{NP})$. Hermitian metrics on the original bundles induce inner products on these spaces.

To summarize, we have the following three sequences of Szegő kernels:

- $\Pi_{NP}^{\mathbb{CP}^m}$ = the orthogonal projection to $H^0(\mathbb{CP}^m, \mathcal{O}(Np))$;
- Π_{NP} = the orthogonal projection to $H^0(\mathbb{CP}^m, \mathcal{O}(Np), NP)$;
- $\Pi_N^{M_P}$ = the orthogonal projection to $H^0(M_P, L_P^N)$.

We note that for the case $P = p\Sigma$, the three sequences coincide. However, if $P \subsetneq p\Sigma$, then the second and third sequences are quite different.

1.3. Moment maps and torus actions. The group $\mathbb{C}^{*m}$ acts on M_P^c and the subgroup $\mathbf{T}^m$ acts in a Hamiltonian fashion. Let us recall the formula for its moment map $\mu_P^c : M_P^c \rightarrow \mathbb{R}^m$, restricted to the open orbit $\mathbb{C}^{*m}$. This moment map is the composition

$$\mu_P^c : M_P^c \subset \mathbb{CP}^{\#P-1} \xrightarrow{\mu_0} \mathbb{R}^{\#P} \xrightarrow{A} \mathbb{R}^m ,$$

where

$$\mu_0([z_1, \dots, z_{\#P}]) = \frac{1}{\|z\|^2}(|z_1|^2, \dots, |z_{\#P}|^2),$$

and A is the linear projection is given by the column vectors $(\alpha^1, \dots, \alpha^{\#P})$. Hence we have

$$\mu_P^c = \frac{1}{\sum_{\alpha \in P} |c_\alpha|^2 |\chi_\alpha|^2} \sum_{\alpha \in P} |c_\alpha|^2 |\chi_\alpha|^2 \alpha . \quad (33)$$

For any c , the image of M_P^c under μ_P^c equals P .

For $\alpha \in P$, we again lift $\chi_\alpha \in H^0(M_P^c, L_P^c)$ to an equivariant function $\hat{\chi}_\alpha^P$ on the circle bundle $X_P^c \rightarrow M_P^c$, and we write

$$\hat{m}_{\alpha(j)}^P := c_{\alpha(j)} \hat{\chi}_{\alpha(j)}^P = \zeta_j \circ \iota_P \quad (34)$$

where $\iota_P : X_P^c \rightarrow S^{2d+1}$ is the lift of the embedding $M_P^c \hookrightarrow \mathbb{CP}^{\#P-1}$. (Of course, $\hat{m}_\alpha^P$ depends on c , which we omit to simplify notation.) We also consider the monomials

$$m_\alpha := c_\alpha \chi_\alpha$$

so that $\hat{m}_\alpha^P$ is the equivariant lift of m_α to X_P^c . In terms of local coordinates (z, θ) on $\pi^{-1}(\mathbb{C}^{*m}) \subset X_P^c$, we have

$$\hat{m}_\alpha^P(z, \theta) = \frac{e^{i\theta} c_\alpha z^\alpha}{\left(\sum_{\beta \in P} |c_\beta|^2 |z^\beta|^2\right)^{1/2}} . \quad (35)$$

Noting that

$$\sum_{\alpha \in P} |\hat{m}_\alpha^P|^2 = \sum_{j=1}^{\#P} |\zeta_j \circ \iota_P|^2 \equiv 1 , \quad (36)$$

we obtain the formula:

$$\mu_P^c(z) = \sum_{\alpha \in P} |\hat{m}_\alpha^P(z)|^2 \alpha . \quad (37)$$

(We write $|\hat{m}_\alpha^P| = |\hat{m}_\alpha^P(z, \theta)|$, since the absolute value is independent of θ .)

In the case where the polytope is the full simplex $p\Sigma$, we shall use the special choice

$$c_\alpha^* = \binom{p}{\alpha}^{\frac{1}{2}} = \left(\frac{p!}{(p+m)!}\right)^{\frac{1}{2}} \|\chi_\alpha\|^{-1} ,$$

so that

$$\mu_{p\Sigma}(z) := \mu_{p\Sigma}^{c^*}(z) = \frac{1}{\sum_{|\alpha| \leq p} \binom{p}{\alpha} |z^\alpha|^2} \sum_{|\alpha| \leq p} \binom{p}{\alpha} |z^\alpha|^2 \alpha = \frac{p}{1 + \sum |z_j|^2} (|z_1|^2, \dots, |z_m|^2) , \quad (38)$$

where the last equality follows by differentiating the identity $(1 + \sum x_j)^p = \sum_{|\alpha| \leq p} \binom{p}{\alpha} x^\alpha$. Note that this choice gives us the scaling formula

$$\mu_{p\Sigma} = p\mu_\Sigma .$$

Furthermore,

$$|\hat{m}_\alpha^{p\Sigma}(z)| = \left[\frac{p!}{(m+p)!} \right]^{\frac{1}{2}} \frac{|\hat{\chi}_\alpha(z)|}{\|\chi_\alpha\|} , \quad (39)$$

where, by abuse of notation, we regard $|\hat{\chi}_\alpha|$ and $|\hat{m}_\alpha^{p\Sigma}|$ as functions on $\mathbb{C}^{*m}$, since they are invariant under the circle action. Recalling that $M_{p\Sigma} \equiv S^{2m+1}/\mathbb{Z}_p$, we have the more precise formula

$$\hat{m}_\alpha^{p\Sigma}(x') = \left[\frac{p!}{(m+p)!} \right]^{\frac{1}{2}} \frac{\hat{\chi}_\alpha(x)}{\|\chi_\alpha\|},$$

where $x' \in M_{p\Sigma}$ is the equivalence class of $x \in S^{2m+1}$, and therefore by (29),

$$\sum_{|\alpha| \leq p} \hat{m}_\alpha^{p\Sigma}(x') \overline{\hat{m}_\alpha^{p\Sigma}(y')} = \langle x, \bar{y} \rangle^p. \quad (40)$$

We now recall a fact on zeros of eigensections which will be useful later on.

PROPOSITION 1.1. *Let $\alpha \in P$. Then for all $z \in M_P$, we have*

$$\chi_\alpha(z) = 0 \Leftrightarrow \mu_P(z) \in \bigcup \{ \bar{F} : F \text{ is a facet}, \alpha \notin \bar{F} \},$$

Proof. We first verify the implication ($\Leftarrow$): Let $z \in \mu_P^{-1}(F)$, $\alpha \notin \bar{F}$. The moment map (37) exhibits $\mu_P(z)$ as a convex combination of $\alpha \in P$. Points on F can however only be convex combinations of lattice points $\alpha \in \bar{F}$. Hence the coefficients of $\alpha \notin \bar{F}$ must vanish. Therefore $\chi_\alpha(z) = 0$.

Since the image under the moment map of the support of $\text{Div } \chi_\alpha$ is a union of facets of P , to prove the reverse implication, it suffices to show that for if F is a facet of P , then $\chi_\alpha \not\equiv 0$ on $\mu_P^{-1}(F)$ whenever $\alpha \in \bar{F}$. But since μ_P is surjective, we see from (37) that this is true whenever α is a vertex of $\bar{F}$. (Otherwise, points of F near α would not be in the image.) Now suppose that $\alpha \in \bar{F}$ is arbitrary. We can write $n_0\alpha = \sum_1^m n_j v_j$, where the v_j are the vertices of $\bar{F}$, $n_j \in \mathbb{N}$ and $\sum n_j = n_0 \geq 1$. Then $\chi_\alpha^{n_0} = \chi_{v_1}^{n_1} \cdots \chi_{v_m}^{n_m} \not\equiv 0$ on $\mu_P^{-1}(F)$. $\square$

Remark: Recall that the polytope P is defined by the equations $\ell_j \geq 0$, where $\ell_j(x) = \langle u_j, x \rangle + a_j$, with u_j the primitive inward normal to the closed facet $\bar{F}_j = \{\ell_j = 0\}$. A more precise statement of Proposition 1.1 is

$$\text{Div } \chi_\alpha = \sum_j \ell_j(\alpha) D_j, \text{ where } D_j = \mu_P^{-1}(\bar{F}_j),$$

for all $\alpha \in \mathbb{Z}^m$. In particular, letting $\alpha = 0$, we see that L_P is the line bundle of the divisor $D_P = \sum a_j D_j$.

We now review how the action of the real torus $\mathbf{T}^m$ lifts from M_P^c to X_P^c and combines with the S^1 action to define a $\mathbf{T}^{m+1}$ action on X_P^c . We use the monomial embedding approach of §1.1.2: Recall that under the monomial embedding

$$\Phi_P^c : \mathbb{C}^{*m} \hookrightarrow M_P^c \hookrightarrow \mathbb{CP}^{\#P-1}, \quad z \mapsto [c_{\alpha(1)} z^{\alpha(1)}, \dots, c_{\alpha(\#P)} z^{\alpha(\#P)}],$$

the $\mathbf{T}^m$ action on $M_P^c \subset \mathbb{CP}^{\#P-1}$ is given by

$$e^{i\varphi} \cdot [\zeta_1, \dots, \zeta_{\#P}] = [e^{i\langle \alpha(1), \varphi \rangle} \zeta_1, \dots, e^{i\langle \alpha(\#P), \varphi \rangle} \zeta_{\#P}]. \quad (41)$$

The action (41) lifts to an action on L_P^{-1} :

$$e^{i\varphi} \cdot \zeta = (e^{i\langle \alpha(1), \varphi \rangle} \zeta_1, \dots, e^{i\langle \alpha(\#P), \varphi \rangle} \zeta_{\#P}). \quad (42)$$

Since the circle bundle $X_P^c \subset S^{2\#P-1}$ is invariant under this action, (42) also gives a lift of the action (41) to X_P^c .

We also have the standard circle action on X_P^c :

$$e^{i\theta} \cdot \zeta = e^{i\theta} \zeta, \quad (43)$$

which commutes with the $\mathbf{T}^m$ -action (42). Combining (42) and (43), we then obtain a $\mathbf{T}^{m+1}$ -action on X_P^c :

$$(e^{i\theta}, e^{i\varphi_1}, \dots, e^{i\varphi_m}) \bullet \zeta = e^{i\theta} (e^{i\varphi} \cdot \zeta). \quad (44)$$

The torus action on X_P^c can be quantized to define an action of the torus as unitary operators on $\mathcal{H}^2(X_P^c) = \bigoplus_{N=0}^{\infty} \mathcal{H}_N^2(X_P^c)$. Specifically, we let $\Xi_1, \dots, \Xi_m$ denote the differential operators on X_P^c generated by the $\mathbf{T}^m$ action:

$$(\Xi_j \hat{S})(\zeta) = \frac{1}{i} \frac{\partial}{\partial \varphi_j} \hat{S}(e^{i\varphi} \cdot \zeta)|_{\varphi=0}, \quad \hat{S} \in \mathcal{C}^\infty(X_P^c). \quad (45)$$

PROPOSITION 1.2. *For $1 \leq j \leq m$,*

- (i) $\Xi_j : \mathcal{H}_N^2(X_P^c) \rightarrow \mathcal{H}_N^2(X_P^c)$;
- (ii) *The lifted monomials $\hat{\chi}_\alpha^P \in \mathcal{H}_N^2(X_P^c)$ satisfy $\Xi_j \hat{\chi}_\alpha^P = \alpha_j \hat{\chi}_\alpha^P$ ($\alpha \in NP$).*

Proof. Item (i) follows from the fact that the $\mathbf{T}^m$ action is holomorphic and commutes with $\frac{\partial}{\partial \theta}$. For the case $N = 1$, (ii) follows immediately from (34) and (42). For $\alpha \in NP$, $N > 1$, we write $\alpha = \beta^1 \cdots \beta^N$ with $\beta^k \in P$, and the conclusion then follows from the first case and the product rule. $\square$

Furthermore, we recall that

$$\frac{\partial}{\partial \theta} : \mathcal{H}_N^2(X_P^c) \rightarrow \mathcal{H}_N^2(X_P^c), \quad \frac{1}{i} \frac{\partial}{\partial \theta} \hat{s}_N = N \hat{s}_N \quad \text{for } \hat{s}_N \in \mathcal{H}_N^2(X_P^c). \quad (46)$$

Remark: The vector fields Ξ_j can be constructed geometrically as follows (see [Gu1]): Let $\xi_j = \frac{\partial}{\partial \varphi_j}$ ($1 \leq j \leq m$) denote the Hamiltonian vector fields generating the $\mathbf{T}^m$ action on M_P^c . There is a natural contact 1-form α on X_P^c determined by the Hermitian connection; a key property of α is that $d\alpha = \pi^*\omega$ (see [Ze, SZ1]). We use α to define the horizontal lifts of the Hamilton vector fields ξ_j :

$$\pi_* \xi_j^h = \xi_j, \quad \alpha(\xi_j^h) = 0.$$

The vector fields Ξ_j are then given by:

$$\Xi_j = \xi_j^h + 2\pi i \langle \mu_P^c \circ \pi, \xi_j^* \rangle \frac{\partial}{\partial \theta} = \xi_j^h + 2\pi i (\mu_P^c \circ \pi)_j \frac{\partial}{\partial \theta}.$$

(Here, $\xi_j^* \in \mathbb{R}^m$ is the element of the Lie algebra of $\mathbf{T}^m$ which acts as ξ_j on M_P .)

2. SZEGÖ KERNELS ON TORIC VARIETIES

As mentioned in the introduction, our analysis of the conditional Szegő kernel is by means of the integral formula (16) relating it to the intrinsic Szegő kernels $\Pi_N^{M_P}$ of M_P . The purpose of this section is to give a special construction of the Szegő kernels of a toric variety which is sufficiently precise for our applications.

Our first step, Proposition 2.1, is to give an exact formula for $\Pi_N^{M_P}$ as the composition of a certain *Toeplitz-Fourier multiplier* denoted $(\mathcal{PQ})^{-1}$ and the pull back of the Fubini-Study kernel under a monomial embedding. The pulled-back Fubini-Study kernel is very simple to analyze and is the *raison d'être* of our method. The Toeplitz-Fourier multiplier requires more work. In Proposition 2.2, we prove that this operator is a $\mathbf{T}^{m+1}$ -invariant *Toeplitz operator* of order m on X_P (modulo a smoothing operator). We apply Proposition 2.2 to obtain the goal of this section, Lemma 2.3, which allows us to use the method of stationary phase to obtain the mass asymptotics of Theorem 2.

2.1. The exact formula. Our exact formula for the Szegő kernel of a toric variety involves two ingredients: the first is the kernel

$$\tilde{\Pi}_1^{M_P}(x, y) := \frac{1}{\delta + 1} \Pi_1^{\mathbb{CP}^\delta}(\iota_P(x), \iota_P(y)) = \langle \iota_P(x), \overline{\iota_P(y)} \rangle = \sum_{\alpha \in P} \hat{m}_\alpha^P(x) \overline{\hat{m}_\alpha^P(y)}, \quad (47)$$

where $\delta = \#P - 1$ and $\iota_P : X_P \rightarrow S^{2\delta+1}$ is the lift of the monomial embedding $M_P \hookrightarrow \mathbb{CP}^\delta$, as described in §1.3. We shall choose the constants $c_\alpha = 1$ for all $\alpha \in P$, so that $\hat{m}_\alpha^P = \hat{\chi}_\alpha^P$. (Except for Proposition 2.1, all our results hold without change for arbitrary $\{c_\alpha\}$.) For simplicity of notation, we shall write

$$\Pi_1^N(x, y) = [\tilde{\Pi}_1^{M_P}(x, y)]^N.$$

We note that $\Pi_1^N(x, x) = 1$.

Second, we introduce two sequences of functions $\mathcal{P}_N$ and $\mathcal{Q}_N$ on NP for each $N = 1, 2, \dots$:

- (i) The ‘partition function’ $\mathcal{P}_N(\alpha) = \#\{(\beta_1, \dots, \beta_N) : \beta_j \in P, \beta_1 + \dots + \beta_N = \alpha\}$, where $\alpha \in NP$. It counts the number of ways of writing a lattice point in NP as a sum of N lattice points each in P .
- (ii) The norming function: $\mathcal{Q}_N(\alpha) := \int_{X_P} |\widehat{\chi}_\alpha^P(x)|^2 d\text{Vol}_{X_P}(x)$.

In our application, we only consider the product

$$\mathcal{P}_N \mathcal{Q}_N(\alpha) = \int_{M_P} \int_{\mathbf{T}} \Pi_1^N(z, e^{i\varphi} z) e^{-i\langle \alpha, \varphi \rangle} d\varphi dV(z).$$

We observe that each sequence of functions can be re-defined as a single function on the homogenized lattice cone $\Lambda_P = \bigcup_{N=1}^{\infty} \widehat{NP}$. By definition, $\widehat{NP} \subset \mathbb{Z}^{m+1}$ is obtained by homogenizing $\alpha \in NP$ to $\widehat{\alpha}^{NP} := (Np - |\alpha|, \alpha_1, \dots, \alpha_m)$. We then define $\mathcal{P}, \mathcal{Q} : \Lambda_P \rightarrow \mathbb{R}_+$ by:

$$\mathcal{P}(\widehat{\alpha}^N) := \mathcal{P}_N(\alpha), \quad \mathcal{Q}(\widehat{\alpha}^N) = \mathcal{Q}_N(\alpha).$$

Since the monomials $\mathcal{Q}_N(\alpha)^{-\frac{1}{2}} \widehat{\chi}_\alpha^P$ form a complete orthonormal basis of $\mathcal{H}^2(X_P)$, they define the eigenvalues of a special kind of operator on $\mathcal{H}^2(X_P)$.

Definition: An operator $\Pi F \Pi$ on $\mathcal{H}^2(X_P)$ will be called a *Toeplitz-Fourier multiplier* if it satisfies the following (equivalent) conditions:

- F may be expressed as a function $F(D)$ of the commuting system of operators $D = (\Xi_1, \dots, \Xi_m, \frac{\partial}{\partial \theta})$.
- Its eigenfunctions are the monomials $\widehat{\chi}_\alpha^P$.

Thus, we put:

$$\mathcal{P}(D) \widehat{\chi}_\alpha^P = \mathcal{P}(\widehat{\alpha}^N) \widehat{\chi}_\alpha^P, \quad \mathcal{Q}(D) \widehat{\chi}_\alpha^P = \mathcal{Q}(\widehat{\alpha}^N) \widehat{\chi}_\alpha^P, \quad \alpha \in NP. \quad (48)$$

The following explicit *factorization formula* exhibits the Szegő kernels $\Pi_N^{M_P}$ as the composition of two simple operators. It is the basis for our analysis of the analytic continuation $\Pi_N^{M_P}$.

PROPOSITION 2.1. *We have:*

$$\Pi_N^{M_P} = (\mathcal{P}\mathcal{Q})^{-1} \Pi_1^N.$$

Proof. First, we have by definition,

$$\Pi_N^{M_P}(x, y) = \sum_{\alpha \in NP} \frac{1}{\mathcal{Q}_N(\alpha)} \widehat{\chi}_\alpha^P(x) \overline{\widehat{\chi}_\alpha^P(y)}.$$

On the other hand, by definition of the partition function, we also have

$$\Pi_1^N(x, y) = \sum_{\alpha \in NP} \mathcal{P}_N(\alpha) \widehat{\chi}_\alpha^P(x) \overline{\widehat{\chi}_\alpha^P(y)}.$$

We note that the N^{th} power of $\Pi_1^{M_P}$ gives the sum over the correct set of exponents but does not have the correct normalizing coefficients. We need to divide each term by $\mathcal{P}_N(\alpha) \mathcal{Q}_N(\alpha)$ to adjust the coefficients. That is just what the Proposition claims. $\square$

Since

$$(\mathcal{P}\mathcal{Q})^{-1} \Pi_1^N = \Pi_N^{M_P} (\mathcal{P}\mathcal{Q})^{-1} \Pi_N^{M_P} \Pi_1^N,$$

we see that $\Pi_N^{M_P}$ is the composition of the Toeplitz-Fourier multiplier $\Pi_N^{M_P} (\mathcal{P}\mathcal{Q})^{-1} \Pi_N^{M_P}$ with the simple kernel Π_1^N .

2.2. Toeplitz operator approach to $\Pi_N^{M_P}$. Our next step is to prove that the Toeplitz-Fourier multiplier $\Pi_N^{M_P}(\mathcal{PQ})^{-1}\Pi_N^{M_P}$ is a more familiar kind of operator, namely a Toeplitz operator.

We begin with some background on symbols. We consider symbols σ of the form $\sigma(z, D_\theta)$, where $\sigma(z, N)$ is a semiclassical symbol. Here, we say $\sigma \in S^k(M \times \mathbb{N})$ is a semiclassical symbol of order k if

$$\sigma_N(z) = \sigma(z, N) = N^k \sum_{j=0}^{l-1} a_j(z) N^{-j} + r_N^l(z), \quad \text{with } \|D_z^n r_N^l(z)\| \leq C_{nl} N^{k-l} \quad (l \geq 1),$$

where $a_j \in \mathcal{C}^\infty(M)$. It is a *smoothing symbol* if $\|D_z^n \sigma_N(z)\| \leq C_n N^{-l}$ for all n, l . The *Toeplitz operator* associated to a symbol σ is the operator $\Pi\sigma(z, D_\theta)\Pi$, where D_θ denotes the symbol of $\frac{\partial}{\partial\theta}$ (the S^1 generator). Its symbol is the polyhomogeneous function on the symplectic cone $\Sigma = \{(x, r\alpha_x) : r > 0\} \subset T^*X$ given by

$$\sigma(z, p_\theta) \sim \sum_{j=0}^{\infty} a_j(z) D_\theta^{k-j}.$$

Since σ commutes with the S^1 action, we have $\Pi\sigma\Pi = \sum_N \Pi_N \sigma_N \Pi_N$.

To simplify notation, we shall write $\Pi_N = \Pi_N^{M_P}$. The goal of this section is to prove that $\Pi_N(\mathcal{PQ})^{-1}\Pi_N$ is a semi-classical Toeplitz operator in the following sense:

PROPOSITION 2.2. *There exists a symbol σ_N of order m with principal symbol equal to 1 and a smoothing operator R_N so that*

$$\Pi_N(\mathcal{PQ})^{-1}\Pi_N = \Pi_N \sigma_N \Pi_N + R_N.$$

As a corollary of Proposition 2.2, we obtain the following lemma, which is the motivation for our approach to $\Pi_{|N_P}$ through $\Pi_N^{M_P}$. It provides an essential ingredient in our derivation of the mass asymptotics.

LEMMA 2.3. *There exists a symbol $\sigma_N \in S^m(M, \mathbb{N})$ with principal symbol equal to 1 and a smoothing operator R_N such that for $t \in \mathbf{T}^m$, we have*

$$\begin{aligned} \int_{X_P} \Pi_N(x, t \cdot x) d\text{Vol}_{X_P}(x) &= \int_{X_P} \sigma_N(x) \Pi_1^N(x, t \cdot x) d\text{Vol}_{X_P}(x) \\ &\quad + \int_{X_P} \int_{X_P} R_N(y, x) \Pi_1^N(x, t \cdot y) d\text{Vol}_{X_P}(x) d\text{Vol}_{X_P}(y). \end{aligned}$$

Proof (assuming Proposition 2.2): Applying $\Pi_1^N = \Pi_N(\mathcal{PQ})\Pi_N$ to the identity of Proposition 2.2, we obtain

$$\Pi_N = (\Pi_N \sigma_N \Pi_N + R_N) \Pi_1^N.$$

Now let T_t denote the translation operator $f(x) \mapsto f(t^{-1} \cdot x)$ on $\mathcal{L}^2(X_P)$. Since $[T_t, \Pi_N] = 0$, we then have

$$T_t \Pi_N = T_t \Pi_N (\sigma_N + R_N) \Pi_1^N = \Pi_N T_t (\sigma_N + R_N) \Pi_1^N.$$

Therefore,

$$\begin{aligned} \int_{X_P} \Pi_N(t^{-1} \cdot x, x) d\text{Vol}_{X_P}(x) &= \text{Trace } T_t \Pi_N = \text{Trace } \Pi_N T_t (\sigma_N + R_N) \Pi_1^N \\ &= \text{Trace } T_t \sigma_N \Pi_1^N + \text{Trace } T_t R_N \Pi_1^N \\ &= \int_{X_P} \sigma_N(t^{-1} \cdot x) \Pi_1^N(t^{-1} \cdot x, x) d\text{Vol}_{X_P}(x) \\ &\quad + \int_{X_P} \int_{X_P} R_N(t^{-1} \cdot x, y) \Pi_1^N(y, x) d\text{Vol}_{X_P}(x) d\text{Vol}_{X_P}(y). \end{aligned}$$

The conclusion follows by the change of variables $x \mapsto t \cdot x$. □

2.2.1. Boutet de Monvel-Sjöstrand parametrix. In proving Proposition 2.2 and other statements about Toeplitz operators, we need some a priori facts about Π^{M_P} which follow from the Boutet de Monvel-Sjöstrand parametrix construction. In this section, we provide the relevant background.

The Boutet de Monvel-Sjöstrand parametrix is a complex oscillatory integral formula for the Szegö kernels of a strictly convex CR manifold modulo smoothing kernels [BS]. It can be used to obtain asymptotics of projection kernels onto holomorphic sections of powers of a positive line bundle over a Kähler manifold (see [Ze, BSZ]). We recall here the basic properties of the Szegö kernel for (M_P, L_P) . Unfortunately, the remainder term is not sufficiently precise for the tunneling theory we have in mind. In the next section, we give a special construction of the Szegö kernels of toric varieties which is sufficiently precise. The construction uses the general properties established in this section.

Now consider any polarized algebraic manifold (M, L) , and assume L is very ample. Choose a basis $\{S_0, \dots, S_d\}$ with $d = \dim H^0(M, L)$ and let Φ denote the associated embedding into projective space. That is, we write $S_j = f_j e_L$ relative to a local frame e_L and put $\Phi(z) = [f_0(z), \dots, f_d(z)]$. Recalling that $L = \Phi^* \mathcal{O}(1)$, we can equip L with the metric $\Phi^* h_{\text{FS}}$. We also give M the metric $\omega = \Phi^* \omega_{\text{FS}}$, and we let Π_N denote the orthogonal projection onto $H^0(M, L^N)$ with respect to these metrics. Also, let $\Pi = \sum_{N=1}^{\infty} \Pi_N$ denote the Szegö kernel.

It was proved by Boutet de Monvel and Sjöstrand [BS] (see also the Appendix to [BG]) that Π is a complex Fourier integral operator of positive type,

$$\Pi \in I_c^0(X \times X, \mathcal{C}) \quad (49)$$

associated to a positive canonical relation $\mathcal{C}$. For definitions and notation concerning complex FIO's we refer to [MS, BS, BG]. The real points of $\mathcal{C}$ form the diagonal $\Delta_{\Sigma \times \Sigma}$ in the square of the symplectic cone

$$\Sigma = \{r\alpha_x : r > 0, x \in X\}, \quad (50)$$

where α is the connection form. We refer to [BG] (see Lemma 4.5 of the Appendix). Moreover, in [BS] a parametrix is constructed for Π , from which it follows (see [Ze, BSZ]) that

$$\Pi_N(x, y) \sim N \int_0^\infty \int_0^{2\pi} e^{N(-i\theta + t\psi(r_\theta x, y))} s(r_\theta x, y, Nt) d\theta dt, \quad (51)$$

where $s(x, y, t) \sim \sum_{k=0}^\infty t^{m-k} s_k(x, y) \in S^m(X \times X \times \mathbb{R}_+)$ is a classical symbol of order m . Here, ' $\sim$ ' means modulo a rapidly decaying term (i.e., a term whose $\mathcal{C}^j$ norms are $O(N^{-k})$ for all j, k).

To describe the phase in (51), we let e_L be a nonvanishing holomorphic section of L over an open $U \subset M$, and consider the analytic extension' $a(z, w)$ of $a(z, z) := a(z) = \|e_L(z)\|_h^{-2}$ in $U \times U$ such that $a(w, z) = \overline{a(z, w)}$ on $U \times U$. Using coordinates (z, θ) for the point $x = e^{i\theta} a(z)^{\frac{1}{2}} e_L(z) \in X$, we have

$$\psi(x_1, x_2) = -1 + e^{i(\theta_1 - \theta_2)} \frac{a(z_1, z_2)}{\sqrt{a(z_1)} \sqrt{a(z_2)}}. \quad (52)$$

Again assuming that the metric ω on M is the pull-back of ω_{FS} , we claim that the phase equals:

$$\psi(x, y) = -1 + \langle \iota(x), \overline{\iota(y)} \rangle, \quad (53)$$

where $\iota : X \rightarrow \mathbb{C}^{2d+1}$ is the lift of Φ given by

$$\iota(z, \theta) = e^{i\theta} \left(\sum |f_j(z)|^2 \right)^{-\frac{1}{2}} (f_0(z), \dots, f_d(z)).$$

To see this, let us recall the Szegö kernel of the hyperplane section bundle $\mathcal{O}(1) \rightarrow \mathbb{CP}^d$ over projective space with the Fubini-Study metric. We take $U = \{z_0 \neq 0\} \approx \mathbb{C}^d$, and we consider the local frame $e = z_0$. Using the local coordinates $[1, z_1, \dots, z_m] \mapsto (z_1, \dots, z_m) \in \mathbb{C}^d$, we then have $a^{\mathbb{CP}^d}(z) = \|e(z)\|^{-2} = 1 + \sum_{j=1}^d |z_j|^2$. Hence $a^{\mathbb{CP}^d}$ has the real-analytic extension

$$a^{\mathbb{CP}^d}(z, w) = 1 + \sum_{j=1}^d z_j \bar{w}_j, \quad (54)$$

and (53) follows. In the case of a toric variety $M_{\mathbb{P}}$, (53) becomes

$$\psi(x, y) = -1 + \sum_{\alpha \in P} \widehat{m}_{\alpha}^P(x) \overline{\widehat{m}_{\alpha}^P(y)}.$$

2.2.2. Complex Fourier integral operators. We now use the theory of complex Fourier integral operators [MS] (or alternatively, Toeplitz operators [BG]) to prove that the Toeplitz Fourier multiplier $\Pi(\mathcal{PQ})^{-1}\Pi$ is a Toeplitz operator of order m modulo a smoothing operator (Proposition 2.2). Recall that a Toeplitz operator in the sense of [BG] is an operator of the form $\Pi A \Pi$ where $A \in \Psi^m(X)$ for some m . Here and in the following, $\Psi^m(X)$ denotes the space of pseudodifferential operators of order m on X . A smoothing Toeplitz operator is a smoothing operator of the form $\Pi R \Pi$.

It follows then that $\Pi_N = T_N \Pi_1^N$ where T_N is a Toeplitz operator. Since it is of some independent interest and requires no extra work, we prove this as a corollary of a general result valid for any polarized algebraic manifold (M, L) . We assume L is very ample, and let $\iota : M \rightarrow \mathbb{CP}^d$ denote the holomorphic embedding induced by $H^0(M, L)$. Let $\Pi_{\iota}(x, y) = \Pi(\iota(x), \iota(y))$ denote the pullback to X of the Fubini-Study Szegő kernel of $\mathbb{CP}^d$. Also, let Π^M or more simply Π denote the Szegő kernel for M with respect to $\iota^* \omega_{\text{FS}}$.

About complex Fourier integral operators A (or Toeplitz operators), all we need to know are the following basic facts:

- $A \in I_c^m(X \times X, \mathcal{C})$ possesses a principal symbol σ_A which is a half-density (times a Maslov factor) along the underlying canonical relation $\mathcal{C}$. (In the Toeplitz case, it is a symplectic spinor.)
- We can compose operators in $I_c^m(X \times X, \mathcal{C})$ on the left and right by elements $B \in \Psi^k(X)$ and $\sigma_{AB} = \sigma_A \sigma_B = \sigma_{BA}$. (The same is true of Toeplitz operators.)
- If $A \in I_c^m(X \times X, \mathcal{C})$ and if $\sigma_A = 0$, then $A \in I_c^{m-1}(X \times X, \mathcal{C})$ (and also for Toeplitz operators).
- Π and Π_{ι} are elliptic in that their symbols are nowhere vanishing. See [BS] for the symbol in the complex FIO sense and [BG, §11] for the symbol in the Toeplitz sense.

LEMMA 2.4. *Let Π denote the Szegő projector associated to the metric $\iota^* \omega_{\text{FS}}$. Then, there exist $A \in \Psi^m(X)$ with $[A, D_{\theta}] = 0$ such that $\Pi \sim \Pi A \Pi \Pi_{\iota} = \Pi A \Pi_{\iota}$ modulo smoothing operators $\Pi R \Pi$.*

Proof. As mentioned above (49), Π is a complex Fourier integral operator associated to a positive canonical relation $\mathcal{C}$, whose real points form the isotropic relation $\Delta_{\Sigma \times \Sigma} \subset T^*X \times T^*X$ (the diagonal). We observe that also $\Pi_{\iota} \in I^*(X \times X, \mathcal{C})$. This follows immediately from the fact, shown by (53), that Π and Π_{ι} are complex Fourier integral distributions with precisely the same phase functions. Since the underlying canonical relation is parametrized by the phase, they both belong to the same class of Fourier integral distributions.

Now, the principal symbol σ_{Π} of Π , viewed as a complex Fourier integral distribution, is a nowhere vanishing $1/2$ -density on $\mathcal{C}$ which is computed in [BS, Prop. 4.8]. Alternatively, viewed as a Toeplitz operator in the sense of [BG] (see Chapter 11), its symbol is an idempotent symplectic spinor. Similarly, the principal symbol $\sigma_{\Pi_{\iota}}$ of Π_{ι} is the pull back under ι of the nowhere vanishing symbol of $\Pi^{\mathbb{CP}^d}$.

By our normalization, Π_{ι} has order $-m$ (since its amplitude is a constant independent of N). We therefore begin by seeking $A_0 \in \Psi^m(X)$ such that $[A_0, D_{\theta}] = 0$ and such that $\Pi - \Pi A_0 \Pi_{\iota}$ is of order -1 . We first find $a_0 \in C^{\infty}(M)$ such that $\sigma_{\Pi} = a_0 \sigma_{\Pi_{\iota}}$ and choose A_0 so that $[A_0, D_{\theta}] = 0$ and so that $\sigma_{A_0} = a_0$. Existence of such an A_0 follows by ellipticity of Π_{ι} and by averaging; see also [BG, Prop. 2.13]. Thus, the principal symbol of order 0 of $\Pi - \Pi A_0 \Pi_{\iota}$ equals zero, i.e. $\Pi - \Pi A_0 \Pi_{\iota} \in I^{-1}(X \times X, \mathcal{C})$. We denote its principal symbol by σ_{-1} . We then seek $A_{-1} \in \Psi^m(X)$ so that $[A_{-1}, D_{\theta}] = 0$, and so that $\Pi - \Pi A_0 \Pi_{\iota} - \Pi A_{-1} D_{\theta}^{-1} \Pi_{\iota} \in I^{-2}(X \times X, \mathcal{C})$. Here, we note that $\Pi D_{\theta} \Pi$ is an elliptic Toeplitz operator; since $\Pi \Pi_{\iota} = \Pi_{\iota}$, the expressions $D_{\theta}^{-1} \Pi_{\iota}$ are well-defined. It suffices to choose $a_{-1} = \sigma_{A_{-1}} \in C^{\infty}(M)$ so that $a_{-1} \sigma_{\Pi_{\iota}} = \sigma_{-1}$. We continue in this way to obtain $a_{-j} \in C^{\infty}(M)$ always using that $\sigma_{\Pi_{\iota}}$ is nowhere vanishing. By a Borel summation argument, we can find $A \in \Psi^m(X)$ with the above commutation properties so that $\Pi A \Pi - \sum_{j=0}^M \Pi A_{-j} D_{\theta}^{-j} \Pi$ is a Toeplitz operator of order $-M - 1$. Then

$$\Pi - \Pi A \Pi_{\iota} \in I^{-\infty}(X \times X, \mathcal{C}).$$

□

We remark that an alternative to the observation that the two complex FIO's have the same phase function is that, by the choice of the Kähler form $\iota^* \omega_{\text{FS}}$, ι is a symplectic (as well as holomorphic) embedding $M \subset \mathbb{CP}^d$. Hence, the pull back operator ι^* carries the class of FIO's in the class of $\Pi^{\mathbb{CP}^d}$ to those in the class of Π .

2.2.3. Semi-classical Toeplitz operators. To complete the proof of Proposition 2.2, we need to relate the complex FIO $\Pi \Pi_l$ to semi-classical Toeplitz operators. The relevant analysis already exists in [Gu2].

We will need the following asymptotic formula for symbols. In the language of Berezin-Toeplitz operators, we are essentially computing the Berezin transform between covariant and contravariant symbols of a Toeplitz operator (see [RT] for background).

LEMMA 2.5. *Let $\sigma_N \sim N^k \sum_{j=0}^{\infty} s_{-j} N^{-j}$ be a semiclassical symbol of order k . Then there exists a complete asymptotic expansion*

$$N^{-m}(\Pi_N \sigma_N \Pi_N)(z, z) = \sum_{j=0}^{l-1} b_{-j}(z) N^{k-j} + r_N^l(z) \quad (l \geq 1),$$

where $b_0 = s_0, b_{-1} = \Delta s_0 + s_{-1}, \dots$, and in general where b_{-j} is a sum of differential operators applied to $s_0, s_{-1}, \dots, s_{-j}$. Also, $\|D_z^n r_N^l(z)\| \leq C_{nl} N^{k-l}$.

Proof. Apply the method of stationary phase using (51)–(52) exactly as in the proof of [Ze, Theorem 1] (which is the case $\sigma = 1$). $\square$

We then have:

LEMMA 2.6. *Let (M, L) be a polarized algebraic manifold as above. Then there exists a symbol $a_N \in S^{-m}(M, \mathbb{N})$ with principal symbol equal to 1 so that*

$$\Pi_1^N = \Pi_N a_N \Pi_N + R_N \quad \text{where} \quad \|R_N\|_{\text{HS}} = O(N^{-k}) \quad \forall k.$$

Here, $\|R_N\|_{\text{HS}}^2 = \text{Trace } R_N^* R_N$.

Proof. By Lemma 2.4, there exists $A \in \Psi^{-m}(X)$ such that $\Pi_l = \Pi A \Pi$. Since $[A, D_\theta] = 0$, it follows by [Gu2] that there exists a symbol a_N with $\|\Pi_1^N - \Pi_N a_N \Pi_N\|_{\text{HS}} = O(N^{-\infty})$. We may determine a_N by using Lemma 2.5. Indeed, we have

$$\Pi_1^N(z, z) = 1 \sim \Pi_N a_N \Pi_N(z, z) \quad (55)$$

modulo functions r_N which tend to zero rapidly in $\mathcal{C}^k(M)$. It follows that $a_0 = 1$ (and then the rest of the coefficients may be determined recursively, e.g. $a_{-1} = 0, a_{-2} = -D_2 a_0$, and so on). $\square$

2.2.4. Completion of the proof of Proposition 2.2. In the toric case, $\Pi_N(\mathcal{PQ})^{-1} \Pi_N$ is the inverse of $\Pi_N(\mathcal{PQ}) \Pi_N = \Pi_1^N$ on $\mathcal{H}^2(X_P)$. From Lemma 2.6 we may write $\Pi_N(\mathcal{PQ}) \Pi_N \sim \Pi_N a_N \Pi_N$ modulo smoothing operators.

We note that we may invert $\Pi a \Pi$ in the class of Toeplitz operators; i.e., there exists a symbol σ_N such that

$$\Pi_N \sigma_N \Pi_N \circ \Pi_N a_N \Pi_N \sim \Pi_N \quad (56)$$

modulo smoothing operators. Such an inverse symbol exists since $a_0 = 1$. The algebraic formalism in which the inverse is calculated is that of $*$ -products of semiclassical symbols. We recall that composition of Toeplitz operators defines a $*$ -product on semiclassical symbols by the formula

$$\Pi_N a_N \Pi_N \circ \Pi_N b_N \Pi_N \sim \Pi_N a_N * b_N \Pi_N. \quad (57)$$

The formula for $*$ may be worked out directly from the parametrix (51) and the inverse can be computed from this formula (see [Gu2, St]).

By (56) we obtain a symbol σ_N (with principal symbol equal to 1) such that

$$\Pi_N \sigma_N \Pi_N \Pi_N(\mathcal{PQ}) \Pi_N \sim \Pi_N.$$

Multiplying both sides by $\Pi_N(\mathcal{PQ})^{-1} \Pi_N$, we conclude the proof. $\square$

Remark: We emphasize that the distinguishing features of the toric case in Proposition 2.2 and Lemma 2.6 are the exact factorization and the fact that the operator $\Pi(\mathcal{PQ}) \Pi$ mediating between Π and Π_l is invertible.

This is due to the fact that sections of L generate the ring $\bigoplus_{N=1}^{\infty} H^0(M, L^N)$ in the toric case. In general they do not and the exact representation $\Pi = B\Pi_L$ in the toric case is only valid modulo smoothing operators.

2.2.5. Alternate proof of Lemma 2.6. One could avoid using the calculus of complex Fourier integral operators or Toeplitz operators in the proof of Proposition 2.2 by further developing the calculus of semi-classical Toeplitz operators directly from the Boutet de Monvel- Sjöstrand parametrix as follows:

We first simplify the expression (51) by using the complex method of stationary phase to eliminate the integrals in the parametrix. The critical point set of the phase $\Phi(\theta, \lambda; x, y) := -i\theta + \lambda\psi(r_\theta x, y)$ is given by

$$\begin{cases} d_\lambda \Phi(\theta, \lambda; x, y) = \psi(r_\theta x, y) = 0 \iff e^{i\theta} \langle \iota(x), \iota(y) \rangle = 1, \\ d_\theta \Phi(\theta, \lambda; x, y) = -i + d_\theta \lambda \psi(r_\theta x, y) = 0 \iff \lambda e^{i\theta} \langle \iota(x), \iota(y) \rangle = 1. \end{cases} \quad (58)$$

It is easy to see by the Schwartz inequality that a real critical point exists if and only if $x = y$, in which case $\theta = 0, \lambda = 1$ and we obtain the familiar expansion along the diagonal. When $x \neq y$ we deform the contour to $|\zeta| = e^\tau$ ($\zeta = e^{i\theta+\tau}$) so that $e^{i\theta+\tau} \langle \iota(x), \iota(y) \rangle = 1$. This is possible as long as $\langle \iota(x), \iota(y) \rangle \neq 0$, as happens near the diagonal, where the parametrix is valid. Because the phase is linear in λ it is clear that the critical point is non-degenerate if and only if $\langle \iota(x), \iota(y) \rangle \neq 0$ and that the Hessian determinant equals $|\langle \iota(x), \iota(y) \rangle|^2$. On the critical set the phase equals $-i\theta + \tau = \log \langle \iota(x), \iota(y) \rangle$, hence we have

$$\Pi_N(x, y) = e^{N \log \langle \iota(x), \iota(y) \rangle} S_N(x, y) + W_N(x, y), \quad (59)$$

where $S_N(x, y) \sim \sum_{k=0}^{\infty} N^{m-k} S_k(x, y)$, and where $W_N(x, y)$ is a smooth uniformly rapidly decaying function. Here, we have absorbed the remainder in the parametrix construction as well as the remainder in the stationary phase expansion of the parametrix in W_N . Note that the first term may be smaller than the second outside a tubular neighborhood of radius $N^{-1/2}$ of the diagonal.

As above, we use Lemma 2.5 to find a symbol a_N so that (55) holds and hence the kernels Π_1^N and $\Pi_N a_N \Pi_N$ agree on the diagonal modulo smoothing symbols. Next, we note that both $\Pi_1^N(x, y)$ and $\Pi_N a_N \Pi_N(x, y)$ are complex oscillatory functions with common phase

$$\Psi(z, w) = \log \langle \iota(x), \iota(y) \rangle. \quad (60)$$

In the case of Π_1^N , this follows from (47). Indeed, we simply have:

$$\Pi_1^N(x, y) = e^{N \log \langle \iota(x), \iota(y) \rangle}. \quad (61)$$

In the case of $\Pi_N a_N \Pi_N$, we apply the method of complex stationary phase to the integral formula

$$\Pi_N a_N \Pi_N(x, y) \sim \int_{M_P} e^{N \Psi(u; x, y)} S_N(x, u) a_N(u) S_N(u, y) dV(u), \quad (62)$$

coming from (59), with

$$\Psi(u; x, y) = \langle \iota(x), \iota(u) \rangle + \langle \iota(u), \iota(y) \rangle.$$

It follows that there exists an amplitude A_N defined near the diagonal such that

$$\Pi_N a_N \Pi_N(x, y) = A_N(x, y) e^{N \log \langle \iota(x), \iota(y) \rangle} + V_N(x, y), \quad (63)$$

where V_N is a new smoothing operator.

Recalling that $X \subset L^*$, we extend A_N to $L^* \times L^*$ so that it is of the form

$$A_N(z, \lambda; w, \lambda') = (\lambda \bar{\lambda}')^N \tilde{A}_N(z, w),$$

where we use a local holomorphic frame to write $x = (z, \lambda)$, $y = (w, \lambda') \in L^*$. We note that $\tilde{A}_N(z, w)$ is holomorphic in z and anti-holomorphic in w near the diagonal. To see this, we first conclude from the construction in [BS] using the $*$ -product that the symbol $s(x, y, t) \sim \sum_{k=0}^{\infty} t^{m-k} s_k(x, y)$ in (51) extends to a symbol on $L^* \times L^*$ that is holomorphic in x and anti-holomorphic in y . It follows by the stationary phase method described above that the same is true for the symbol $S_N(x, y)$ in (59), and then by (63) that the same is also true for A_N as claimed. (Note that in terms of coordinates (z, θ) on X , the function $z \mapsto A_N(z, \theta; w, \theta') = e^{iN(\theta-\theta')} \sqrt{a(z)a(w)} \tilde{A}_N(z, w)$ is not holomorphic in z .)

The amplitude of $\Pi_1^N - \Pi_N a_N \Pi_N$ becomes $(\lambda \bar{\lambda}')^N (1 - \tilde{A}_N(z, w))$, where we can write $1 - \tilde{A}_N \sim B_0 + B_1 N^{-1} + B_2 N^{-2} + \dots$. Since $\Pi_1^N - \Pi_N a_N \Pi_N(z, z) = O(N^{-\infty})$ by our choice of a_N above, it follows that

$B_j(z, z) = 0$ (for all j). Since $B_j(z, w)$ vanishes on the diagonal and is holomorphic in z and anti-holomorphic in w , it must be identically 0. Hence, $\Pi_1^N - \Pi_N a_N \Pi_N(x, y)$ is a smoothing operator. $\square$

3. MASS ASYMPTOTICS

In this section, we prove a precise asymptotic formula for the conditional Szegő kernel on the diagonal (Proposition 3.1), which yields the mass asymptotics of Theorem 2.

First, we discuss formula (8) for the expected mass density. Let P denote a Delzant polytope in $\mathbb{R}^m$. Recalling the definition (5) of the conditional probability measure $\gamma_{P|P}$ on the space $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ of polynomials with Newton polytope P , we see that the expected value of the mass density with respect to $\gamma_{P|P}$ is given by:

$$\mathbf{E}_{|P}(|f(z)|_{\text{FS}}^2) = \sum_{\alpha, \beta \in P} \frac{\mathbf{E}(\lambda_\alpha \bar{\lambda}_\beta) \chi_\alpha(z) \overline{\chi_\beta(z)}}{\|\chi_\alpha\| \|\chi_\beta\| (1 + \|z\|^2)^{p/2}}.$$

Since the λ_α are independent complex random variables with variance 1 (i.e., $\mathbf{E}_{|P}(\lambda_\alpha \bar{\lambda}_\beta) = \delta_\alpha^\beta$), we have:

$$\mathbf{E}_{|P}(|f(z)|_{\text{FS}}^2) = \sum_{\alpha \in P} \frac{|\chi_\alpha(z)|_{\text{FS}}^2}{\|\chi_\alpha\|^2} = \Pi_{|P}(z, z). \quad (64)$$

It then follows by expressing the Gaussian in spherical coordinates that

$$\mathbf{E}_{\nu_P}(|f(z)|_{\text{FS}}^2) = \frac{1}{\#P} \mathbf{E}_{|P}(|f(z)|_{\text{FS}}^2) = \frac{1}{\#P} \Pi_{|P}(z, z).$$

Replacing P with NP , we obtain formula (8).

The number of lattice points in the polytope NP is given by the Riemann-Roch formula (see [Fu]):

$$\begin{aligned} \#(NP) &= \dim H^0(\mathbb{CP}^m, \mathcal{O}(NP), NP) = \dim H^0(M_P, L_P^N) \\ &= \chi(M_P, L_P^N) = \sum_{k=0}^m \deg[c_1(L_P)^k \cup \text{Todd}_{m-k}(M_P)] \frac{N^k}{k!} \\ &= \text{Vol}(P)N^m + \cdots + \deg \text{Todd}_m(M_P). \end{aligned}$$

Here, we used the Kodaira vanishing theorem and the fact that

$$\text{Vol}(P) = \text{Vol}(M_P) = \int_{M_P} \frac{1}{m!} \omega_P = \frac{1}{m!} \deg[c_1(L_P)^m].$$

(This immediately yields Kouchnirenko's Theorem, since $\deg[c_1(L)^m]$ equals the number of points in the intersection of m generic divisors of a very ample line bundle L ; see [At].) The holomorphic Euler characteristic $\chi(M_P, L_P^N)$ is also known as the Hilbert polynomial of the polarized manifold (M_P, L_P) . Combining (8) and the Riemann-Roch formula, we obtain

$$\mathbf{E}_{\nu_{NP}}(|f(z)|_{\text{FS}}^2) = \frac{1}{\chi(M_P, L_P^N)} \Pi_{|NP}(z, z). \quad (65)$$

The mass asymptotics of Theorem 2 is an immediate consequence of (65) and the following asymptotic expansion of the conditional Szegő kernel on the diagonal:

PROPOSITION 3.1. *Suppose that P is a Delzant polytope in $\mathbb{R}^m$. Then:*

i) *For z in the classically allowed region $\mathcal{A}_P$, we have*

$$\Pi_{|NP}(z, z) = \prod_{j=1}^m (Np + j) + R_N(z), \quad \|R_N\|_{C^k(K)} = O(e^{-\lambda_K N}) \quad \forall k,$$

for all compact $K \subset \mathcal{A}_P$, where $\lambda_K > 0$.

ii) *On each open forbidden region $\mathcal{R}_F^\circ$,*

$$\Pi_{|NP}(z, z) = N^{\frac{m+r}{2}} e^{-Nb(z)} [c_0^F(z) + c_1^F(z)N^{-1} + \cdots + c_k^F(z)N^{-k} + R_k^F(z)],$$

where $r = \dim F$ and

- (a) $c_j^F \in \mathcal{C}^\infty(\mathcal{R}_F^\circ)$ and $c_0^F > 0$ on $\mathcal{R}_F^\circ$;
- (b) $\|R_k^F\|_{\mathcal{C}^j(K)} = O(N^{-k-1})$, for all compact $K \subset \mathcal{R}_F^\circ$ and for all j, k ;
- (c) $b > 0$ on $\mathbb{C}^{*m} \setminus \overline{\mathcal{A}_P}$;
- (d) b is given by formula (10);
- (e) $b \in \mathcal{C}_{\mathbb{R}}^1(\mathbb{C}^{*m})$ (with $b = 0$ on $\mathcal{A}_P$), and b is $\mathcal{C}^\infty$ on each closed region $\overline{\mathcal{R}_F}$.

We remark that the asymptotics of Theorem 3.1 are uniform away from the transition points only. To take care of the transition points, we shall also prove the following local uniform convergence result on all of $\mathbb{C}^{*m}$:

PROPOSITION 3.2. *Let P be a Delzant polytope. Then*

$$\frac{1}{N} \log \Pi_{|NP}(z, z) \rightarrow -b(z)$$

*uniformly on all compact subsets of $\mathbb{C}^{*m}$.*

This section is devoted to the proofs of Propositions 3.1 and 3.2. We begin with part (ii) of the former, which is the major part of the section. The proof begins by using Lemma 2.3 to obtain an oscillatory integral formula for the conditional Szegő kernel $\Pi_{|NP}(z, z)$. We then observe that the integral is rapidly decaying when z is in the classically forbidden region, while for z in the allowed region, the phase is of positive type with a nondegenerate critical manifold, and hence $\Pi_{|NP}(z, z)$ has an asymptotic expansion. In §3.2, we consider z in the forbidden region and we seek deformations of the contour of integration so that the (analytic continuation of) the phase picks up critical points. In §3.2.1, we formulate necessary geometric conditions for the existence of critical points where the phase has maximal real part and we show that there is a contour for which these conditions are satisfied. In §§3.2.2–3.2.3, we show that the conditions are sufficient and, with the added assumption that z is not a transition point, the resulting critical points are nondegenerate (Lemma 3.9). We complete the proof of Proposition 3.1(ii) in §§3.2.4–3.2.5. In §3.2.5, we use 3.1(ii) to prove Proposition 3.2, which we then use to prove 3.1(i) in §3.2.6.

We start with a simple integral formula for the conditional Szegő kernel:

LEMMA 3.3. *For $y_1, y_2 \in \mathbb{C}^{*m}$, we have*

$$\Pi_{|NP}(y_1, y_2) = \int_{X_P} \int_{\mathbf{T}^m} \Pi_{NP}^{\mathbb{C}P^m}(t \cdot y_1, y_2) \Pi_N^{M_P}(x, t \cdot x) d\text{Vol}_{\mathbf{T}^m}(t) d\text{Vol}_{X_P}(x).$$

Proof. We consider the lattice sums

$$\chi_{NP}(t) := \sum_{\alpha \in NP} t^\alpha, \quad t \in \mathbf{T}^m. \quad (66)$$

We easily see that

$$\Pi_{|NP}(y_1, y_2) = \int_{\mathbf{T}^m} \Pi_{NP}^{\mathbb{C}P^m}(t \cdot y_1, y_2) \chi_{NP}(\bar{t}) d\text{Vol}_{\mathbf{T}^m}(t). \quad (67)$$

Recalling (32), we also have

$$\int_{X_P} \Pi_N^{M_P}(x, t \cdot x) d\text{Vol}_{X_P}(x) = \int_{X_P} \sum_{\alpha \in NP} \frac{\bar{t}^\alpha}{\|\widehat{\chi}_\alpha^P\|_{X_P}^2} |\widehat{\chi}_\alpha^P(x)|^2 d\text{Vol}_{X_P}(x) = \chi_{NP}(\bar{t}). \quad (68)$$

The conclusion follows immediately from (67)–(68). $\square$

By Lemmas 2.3 and 3.3, we have for $z \in \mathbb{C}^{*m}$,

$$\begin{aligned} \Pi_{|NP}(z, z) &= \frac{1}{(2\pi)^m} \int_{M_P} \int_{\mathbf{T}^m} \Pi_{NP}^{\mathbb{C}P^m}(e^{i\varphi} \cdot z, z) \Pi_N^{M_P}(w, e^{i\varphi} \cdot w) d\varphi d\text{Vol}_{M_P}(w) \\ &= K_N(z) + S_N(z), \end{aligned} \quad (69)$$

where

$$K_N(z) = \frac{1}{(2\pi)^m} \int_{M_P} \int_{\mathbf{T}^m} \Pi_{Np}^{\mathbb{CP}^m}(e^{i\varphi} \cdot z, z) \sigma_N(w) \tilde{\Pi}_1^{M_N}(w, e^{i\varphi} \cdot w) d\varphi dw, \quad (70)$$

$$S_N(z) = \frac{1}{(2\pi)^m} \int_{M_P} \int_{M_P} \int_{\mathbf{T}^m} \Pi_{Np}^{\mathbb{CP}^m}(e^{i\varphi} \cdot z, z) R_N(w, \eta) \tilde{\Pi}_1^{M_N}(\eta, e^{i\varphi} \cdot w) d\varphi dw d\eta, \quad (71)$$

where σ_N , R_N are as in Lemma 2.3, and $\tilde{\Pi}_1^{M_N}$ is given by (47). (The above are really functions of $x = (z, \theta)$, but since they depend only on z , we dropped the variable θ . In (71), we replaced integrals over X_P with integrals over M_P by setting $R_N(w, \eta) = \int_0^{2\pi} \int_0^{2\pi} R_N((w, \theta_1), (\eta, \theta_2)) e^{iN(\theta_1 - \theta_2)} \frac{d\theta_1}{2\pi} \frac{d\theta_2}{2\pi}$.)

By (40), we have

$$\Pi_{Np}^{\mathbb{CP}^m}(x, y) = \frac{(Np+m)!}{(Np)!} \langle x, y \rangle^{pN} = \frac{(Np+m)!}{(Np)!} \left(\sum_{|\alpha| \leq p} \hat{m}_\alpha^{p\Sigma}(x) \overline{\hat{m}_\alpha^{p\Sigma}(y)} \right)^N.$$

Recalling (47), we can then rewrite (70):

$$K_N(z) = \frac{1}{(2\pi)^m} \int_{M_P} \int_{\mathbf{T}^m} e^{N\Psi(\varphi, w; z)} a_N(w) d\varphi d\text{Vol}_{M_P}(w), \quad (72)$$

where the phase is given by

$$\Psi(\varphi, w; z) = \log \sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2 + \log \sum_{|\alpha| \leq p} e^{i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^{p\Sigma}(z)|^2, \quad (73)$$

and $a_N(w) = \frac{(Np+m)!}{(Np)!} \sigma_N(w)$ is a symbol of order $2m$.

We fix $z \in \mathbb{C}^{*m}$. It follows from the triangle inequality and (36) that

$$\begin{aligned} \Re \Psi(\varphi, w; z) &= \log \left| \sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2 \right| + \log \left| \sum_{|\alpha| \leq p} e^{i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^{p\Sigma}(z)|^2 \right| \\ &\leq \Re \Psi(0, w; z) = 0, \end{aligned} \quad (74)$$

with equality if and only if $\langle \alpha, \varphi \rangle = \langle 0, \varphi \rangle = 0$ for all $\alpha \in p\Sigma$, or equivalently, if $\varphi = 0$. This says that Ψ is a phase function of positive type, and $\Re \Psi(\varphi, w) = 0$ if and only if $\varphi = 0$. Hence we need consider only critical points with $\varphi = 0$.

Suppose that $w = e^{i\theta} e^\rho \in \mathbb{C}^{*m}$. We note that Ψ is independent of θ ; hence the critical set is $\mathbf{T}^m$ -invariant. So we must find the set of w where

$$D_\varphi \Psi(0, w) = 0, \quad D_\rho \Psi(0, w) = 0.$$

We have

$$iD_\varphi \Psi = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2 \alpha}{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2} - \frac{\sum_{\alpha \in p\Sigma} e^{i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^{p\Sigma}(z)|^2 \alpha}{\sum_{\alpha \in p\Sigma} e^{i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^{p\Sigma}(z)|^2}. \quad (75)$$

Hence the D_φ equation gives

$$\frac{1}{p} \mu_P(w) = \mu_\Sigma(z). \quad (76)$$

Note that

$$D_\rho \Psi(0, w) = D_\rho 0 \equiv 0$$

automatically by (60). However, we will need to compute $D_\rho \Psi$. To do this, we note that

$$\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2 = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |m_\alpha(w)|^2}{\sum_{\alpha \in P} |m_\alpha(w)|^2}.$$

Recalling that $w = e^{i\theta} e^\rho$, we then obtain

$$\frac{1}{2} D_\rho \Psi = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2 \alpha}{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\hat{m}_\alpha^P(w)|^2} - \mu_P(w). \quad (77)$$

Let $\mathcal{C}_\Psi^0 \subset \mathbf{T}^m \times M_P$ denote the set of critical points of Ψ where $\Re \Psi = 0$. By (74) and (76), we have:

$$\mathcal{C}_\Psi^0 = \{\varphi = 0, w \in \mu_P^{-1}(\mu_\Sigma(z))\} \simeq \mathbf{T}^m. \quad (78)$$

Equation (78) leads to the important conclusion:

There are no critical points with real part equal to 0 unless $z \in (\mu_\Sigma)^{-1}(\frac{1}{p}P)$.

As we stated in the introduction, we refer to the region $\mathcal{A}_P = (\mu_\Sigma)^{-1}(\frac{1}{p}P^\circ)$ as the *classically allowed* region, and its complement in $\mathbb{C}^{*m}$ as the *classically forbidden* region. We now break up our problem into these two regions.

3.1. The classically allowed region. In this section, we illustrate our approach by showing that

$$\Pi_{|NP}(z, z) \sim N^m [c_0(z) + c_1(z)N^{-1} + c_2(z)N^{-2} + \dots], \quad \text{for } z \in \mathcal{A}_P. \quad (79)$$

In §3.2, we will derive our exponentially decaying, asymptotic expansion of $\Pi_{|NP}(z, z)$ on the forbidden region, which we will use to obtain the precise formula of Proposition 3.1(i) on the allowed region. (Although we do not need (79) in the derivation of our precise formula in §3.2, we shall use the methods and Lemma 3.4 from this section in our proof.)

Suppose that $z \in \mathcal{A}_P = (\mu_\Sigma)^{-1}(\frac{1}{p}P^\circ)$ is in the classically allowed region. Since the phase function is of positive type and equals zero on the critical set, in order to apply the method of stationary phase it suffices to show that the Hessian of $\Psi(\varphi, \rho)$ is nondegenerate on the critical set. We write the Hessian as a block matrix:

$$D^2\Psi|_{(0,\rho)} = \begin{pmatrix} A & B \\ B^t & C \end{pmatrix},$$

where

$$A_{jk} = \frac{\partial^2 \Psi}{\partial \varphi_j \partial \varphi_k}(0, \rho), \quad B_{jk} = \frac{\partial^2 \Psi}{\partial \varphi_j \partial \rho_k}(0, \rho), \quad C_{jk} = \frac{\partial^2 \Psi}{\partial \rho_j \partial \rho_k}(0, \rho).$$

Differentiating (75) and recalling that $\mu_P(w) = \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \alpha$, we obtain

$$A = \mu_P(w) \otimes \mu_P(w) - \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \alpha \otimes \alpha + \mu_{p\Sigma}(z) \otimes \mu_{p\Sigma}(z) - \sum_{|\alpha| \leq p} |\widehat{m}_\alpha^{p\Sigma}(z)|^2 \alpha \otimes \alpha.$$

(For a vector $v = (v_1, \dots, v_m) \in \mathbb{R}^m$, we identify $v \otimes v$ with the symmetric matrix $(v_j v_k)_{1 \leq j, k \leq m}$.)

Thus on $\mathcal{C}_\Psi^0$, we have by (76),

$$A = 2p^2 \mu_\Sigma(z) \otimes \mu_\Sigma(z) - \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \alpha \otimes \alpha - \sum_{|\alpha| \leq p} |\widehat{m}_\alpha^{p\Sigma}(z)|^2 \alpha \otimes \alpha. \quad (80)$$

On the other hand, differentiating (77), we obtain

$$B = -2i \left(\sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \alpha \otimes \alpha - \mu_P(w) \otimes \mu_P(w) \right). \quad (81)$$

Finally, since $\Psi(0, \rho) \equiv 0$, we have

$$C = 0.$$

Therefore

$$\det D^2\Psi|_{(0,\rho)} = (\det B)^2. \quad (82)$$

In order to apply the method of stationary phase to conclude that the integral (72) admits a complete asymptotic expansion in powers of N^{-1} , we must show that $\det D^2\Psi|_{(0,\rho)} \neq 0$. This follows from (82) and the following fact:

LEMMA 3.4. *The real symmetric matrix*

$$S_w := \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \alpha \otimes \alpha - \mu_P(w) \otimes \mu_P(w)$$

*is strictly positive definite for all $w \in \mathbb{C}^{*m}$. Furthermore, if $\mu_P(w)$ lies in a face F of P , then S_w is semi-positive and its 0 eigenspace is $T_F^\perp$; in particular, S_w is positive definite on T_F .*

Proof. We must show that $(\lambda \otimes \lambda, S_w) > 0$ for $\lambda \in (\mathbb{R}^m)' \setminus \{0\}$. Consider the vectors $u, v \in \mathbb{R}^{\#P}$ given by $u_\alpha = |\widehat{m}_\alpha^P(w)|$, $v_\alpha = |\widehat{m}_\alpha^P(w)|\lambda(\alpha)$. Since $\|u\|^2 = \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 = 1$ and $\mu_P(w) = \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \alpha$, we have

$$\begin{aligned} (\lambda \otimes \lambda, S_w) &= \sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \lambda(\alpha)^2 - \left(\sum_{\alpha \in P} |\widehat{m}_\alpha^P(w)|^2 \lambda(\alpha) \right)^2 \\ &= \|v\|^2 - \langle u, v \rangle^2 = \|u\|^2 \|v\|^2 - \langle u, v \rangle^2 \geq 0. \end{aligned} \quad (83)$$

If $w \in \mathbb{C}^{*m}$, then the Cauchy-Schwartz inequality in (83) is strict since $|\widehat{m}_\alpha^P(w)| \neq 0$ for all $\alpha \in P \cap \mathbb{Z}$ and $\lambda(\alpha)$ is not constant on $P \cap \mathbb{Z}$. On the other hand, if $\mu_P(w)$ lies in a face F , then by Proposition 1.1, $|\widehat{m}_\alpha^P(w)| \neq 0 \Leftrightarrow \alpha \in \bar{F}$, and hence

$$(\lambda \otimes \lambda, S_w) = 0 \Leftrightarrow \lambda|_F \equiv \text{const.} \Leftrightarrow \lambda \perp T_F.$$

□

The asymptotic expansion (79) for z in the classically allowed region $\mathcal{A}_P$ now follows from the method of stationary phase ([Hö, Theorem 7.7.5]) applied to the complex oscillatory integral (72) and the fact that $S_N(z)$ is rapidly decaying. To be precise, the asymptotics of (72) are determined by the component of the critical point set of the phase on which $\Re \Psi$ is maximal. Note that $\Re \Psi$ and $\Im \Psi$ are constant on components of the critical set, and that on its maximal component our phase satisfies $\Re \Psi = 0$. The critical point set is a manifold and we have just shown that the phase is non-degenerate in the normal directions. Although ([Hö, Theorem 7.7.5]) assumes the critical point with $\Re \Psi = 0$ is isolated, the proof in [Hö] can be modified, precisely as in the real case, to apply to phases with non-degenerate critical manifolds.

Hence the expansion follows from the complex stationary phase method for phases of positive type with non-degenerate critical manifolds. Since the critical manifold is of (real) codimension $2m$ and the amplitude a_N in (72) is of order $2m$, it follows that the leading term of the expansion contains $N^{2m-m} = N^m$, as claimed.

3.2. The classically forbidden region. For z in this region of $\mathbb{C}^{*m}$, there are no critical points of the phase with real part 0, and the integral is rapidly decaying by (74). Perhaps surprisingly, this implies that $\partial \bar{\partial} \log \Pi_{NP}(z, z)$ will contribute here to the distribution of zeros.

To determine the exponential decay rate of the integral (72), we shall deform the contour of integration to pick up critical points with maximal real part along the contour. To accomplish this, we first complexify the real torus $\mathbf{T}^m$ to $\mathbb{C}^{*m}$ with variables $\zeta_j = e^{\tau_j + i\varphi_j}$. The integral (72) may be written in terms of these variables as

$$K_N(z) = \frac{1}{(2\pi i)^m} \int_{M_P} \int_{\mathbf{T}^m} e^{N\Psi_{\mathbb{C}}(\zeta, w; z)} \sigma_N(w) \prod_{j=1}^m \frac{d\zeta_j}{\zeta_j} d\text{Vol}_{M_P}(w), \quad (84)$$

where

$$\begin{aligned} \Psi_{\mathbb{C}}(\zeta, w; z) &= \log \sum_{\alpha \in P} \zeta^{-\alpha} |\widehat{m}_\alpha^P(w)|^2 + \log \sum_{|\alpha| \leq p} \zeta^\alpha |\widehat{m}_\alpha^{p\Sigma}(z)|^2 \\ &= \log \sum_{\alpha \in P} e^{-\langle \alpha, \tau + i\varphi \rangle} |\widehat{m}_\alpha^P(w)|^2 + \log \sum_{|\alpha| \leq p} e^{\langle \alpha, \tau + i\varphi \rangle} |\widehat{m}_\alpha^{p\Sigma}(z)|^2. \end{aligned} \quad (85)$$

Since the integrand is holomorphic in $\zeta \in \mathbb{C}^{*m}$, we can deform the contours in (84) and instead integrate over

$$\int_{M_P} \int_{|\zeta_1|=\tau_1, \dots, |\zeta_m|=\tau_m} \cdot \quad (86)$$

We therefore look for complex critical points of $\Psi_{\mathbb{C}}$. We note that as before,

$$\Re \Psi_{\mathbb{C}}(e^{\tau + i\varphi}, w) < \Re \Psi_{\mathbb{C}}(e^\tau, w), \quad \text{for } \varphi \neq 0. \quad (87)$$

Thus the critical points Ψ with maximal real values on the contour (86) are those with $\varphi = 0$.

We now write $\Psi_{\mathbb{C}}(\tau, \varphi, w) = \Psi_{\mathbb{C}}(e^{\tau+i\varphi}, w; z)$. The complexification of (75) becomes

$$\begin{aligned} iD_{\varphi}\Psi_{\mathbb{C}} &= \frac{\sum_{\alpha \in P} e^{-\langle \alpha, \tau+i\varphi \rangle} |\widehat{m}_{\alpha}^P(w)|^2 \alpha}{\sum_{\alpha \in P} e^{-\langle \alpha, \tau+i\varphi \rangle} |\widehat{m}_{\alpha}^P(w)|^2} - \frac{\sum_{\alpha \in p\Sigma} e^{\langle \alpha, \tau+i\varphi \rangle} |\widehat{m}_{\alpha}^{p\Sigma}(z)|^2 \alpha}{\sum_{\alpha \in p\Sigma} e^{\langle \alpha, \tau+i\varphi \rangle} |\widehat{m}_{\alpha}^{p\Sigma}(z)|^2} \\ &= \frac{\sum_{\alpha \in P} e^{-\langle \alpha, \tau+i\varphi \rangle} |m_{\alpha}(w)|^2 \alpha}{\sum_{\alpha \in P} e^{-\langle \alpha, \tau+i\varphi \rangle} |m_{\alpha}(w)|^2} - \frac{\sum_{\alpha \in p\Sigma} e^{\langle \alpha, \tau+i\varphi \rangle} |m_{\alpha}(z)|^2 \alpha}{\sum_{\alpha \in p\Sigma} e^{\langle \alpha, \tau+i\varphi \rangle} |m_{\alpha}(z)|^2} \\ &= \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2 \alpha}{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2} - \frac{\sum_{\alpha \in p\Sigma} e^{i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^{p\Sigma}(e^{\tau/2} \cdot z)|^2 \alpha}{\sum_{\alpha \in p\Sigma} e^{i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^{p\Sigma}(e^{\tau/2} \cdot z)|^2}, \end{aligned} \quad (88)$$

for $w \in \mathbb{C}^{*m}$. Hence by continuity, we see that

$$iD_{\varphi}\Psi_{\mathbb{C}} = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2 \alpha}{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2} - \frac{\sum_{\alpha \in p\Sigma} e^{i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^{p\Sigma}(e^{\tau/2} \cdot z)|^2 \alpha}{\sum_{\alpha \in p\Sigma} e^{i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^{p\Sigma}(e^{\tau/2} \cdot z)|^2}, \quad (89)$$

for all $w \in M_P$. In particular,

$$iD_{\varphi}\Psi_{\mathbb{C}}(\tau, 0, w) = \mu_P(e^{-\tau/2} \cdot w) - \mu_{p\Sigma}(e^{\tau/2} \cdot z). \quad (90)$$

Hence the equation $D_{\varphi}\Psi_{\mathbb{C}}(\tau, 0, w) = 0$ is equivalent to

$$\frac{1}{p} \mu_P(e^{-\tau/2} \cdot w) = \mu_{\Sigma}(e^{\tau/2} \cdot z). \quad (91)$$

To compute the w derivative, we first consider the case where $w = e^{\rho+i\theta} \in \mathbb{C}^{*m}$. Recalling that

$$\sum_{\alpha \in P} e^{-\langle \alpha, \tau+i\varphi \rangle} |\widehat{m}_{\alpha}^P(w)|^2 = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |m_{\alpha}(e^{-\tau/2} \cdot w)|^2}{\sum_{\alpha \in P} |m_{\alpha}(w)|^2} = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |m_{\alpha}(e^{-\tau/2+\rho})|^2}{\sum_{\alpha \in P} |m_{\alpha}(e^{\rho})|^2},$$

we obtain

$$\frac{1}{2} D_{\rho}\Psi_{\mathbb{C}} = \frac{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2 \alpha}{\sum_{\alpha \in P} e^{-i\langle \alpha, \varphi \rangle} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2} - \mu_P(w). \quad (92)$$

Setting $\varphi = 0$,

$$D_{\rho}\Psi_{\mathbb{C}}(\tau, 0, w) = \mu_P(e^{-\tau/2} w) - \mu_P(w). \quad (93)$$

Hence the equation $D_{\rho}\Psi_{\mathbb{C}}|_{\varphi=0} = 0$ gives

$$\mu_P(e^{-\tau/2} \cdot w) = \mu_P(w). \quad (94)$$

We must show that (94) also holds for critical points $w \notin \mathbb{C}^{*m}$. (These are points with $\mu_P(w) \in \partial P$.) To do this we introduce the function

$$H_P(\tau, w) := \frac{\sum_{\alpha \in P} e^{-\langle \alpha, \tau \rangle} |m_{\alpha}(w)|^2}{\sum_{\alpha \in P} |m_{\alpha}(w)|^2} = \sum_{\alpha \in P} e^{-\langle \alpha, \tau \rangle} |\widehat{m}_{\alpha}^P(w)|^2. \quad (95)$$

Note that $H_P(\tau, \cdot)$ is a well-defined function on M_P that factors through the moment map μ_P . We also have

$$H_{p\Sigma}(\tau, z) = H_{\Sigma}(\tau, z)^p = \left[\frac{1 + \|e^{-\tau/2} \cdot z\|^2}{1 + \|z\|^2} \right]^p. \quad (96)$$

We have

$$\Psi_{\mathbb{C}}(\tau, 0, w; z) = \log H_P(\tau, w) + \log H_{p\Sigma}(-\tau, z), \quad (97)$$

and hence

$$D_w \Psi_{\mathbb{C}}(\tau, 0, w; z) = D_w \log H_P(\tau, w).$$

Furthermore the vector fields

$$X_j := \frac{\partial}{\partial \rho_j} = \Re(w_j \frac{\partial}{\partial w_j}) \quad (98)$$

extend to M_P as the generators of the $\mathbb{R}_+^m$ action (induced by the $\mathbb{C}^{*m}$ action). Hence (93) is equivalent to the equation

$$(X_1, \dots, X_m) \log H_P(\tau, w) = \mu_P(e^{-\tau/2} w) - \mu_P(w), \quad (99)$$

which is invariantly defined on all of M_P .

Thus we have seen that critical points of $H_P(\tau, \cdot)$ must satisfy (94). We shall show later that (94) is also a sufficient condition for w to be a critical point of $H_P(\tau, \cdot)$, and hence (91) and (94) give necessary and sufficient conditions for $(0, w)$ to be a critical point of $\Psi_{\mathbb{C}}(\tau, \cdot, \cdot)$.

The following fact is useful in describing the critical set.

LEMMA 3.5. *Let $\tau \in \mathbb{R}^m$, $w \in M_P$. Then the following are equivalent:*

- a) $\mu_P(e^\tau \cdot w) = \mu_P(w)$;
- b) $e^{r\tau} \cdot w = w$, for all $r \in \mathbb{R}$;
- c) τ is perpendicular to T_{F^w} , where F^w is the face of P containing $\mu_P(w)$.

Proof. Let $w = (w_\alpha)_{\alpha \in P} \in M_P \subset \mathbb{CP}^{\#P-1}$. By Proposition 1.1, we see that $w_\alpha = 0$ if and only if $\alpha \notin \overline{F^w}$, and thus for any $t \in \mathbb{C}^{*m}$,

$$t \cdot w = w \Leftrightarrow \exists c \text{ such that } t^\alpha w_\alpha = c w_\alpha \quad \forall \alpha \in P \Leftrightarrow t^\alpha = c \quad \forall \alpha \in \overline{F^w} \cap \mathbb{Z}^m. \quad (100)$$

Suppose (a) holds. Then there exists $e^{i\varphi} \in \mathbf{T}^m$ such that $e^{\tau+i\varphi} \cdot w = w$. Hence by (100), $e^{\langle \tau+i\varphi, \alpha \rangle} = c$ and thus $e^{\langle r\tau, \alpha \rangle} = |c|^r$, for all $\alpha \in \overline{F^w} \cap \mathbb{Z}^m$. Therefore (b) holds.

Furthermore, it follows from (100) that (b) is equivalent to

$$\langle \tau, \alpha - \alpha' \rangle = 0 \quad \forall \alpha, \alpha' \in \overline{F^w} \cap \mathbb{Z}^m, \quad (101)$$

which is a restatement of (c). $\square$

Remark: Let $\tau \in \mathbb{R}^m$. It follows from Lemma 3.5 that the set A_τ of points in M_P satisfying (94) consists of the inverse image under μ_P of the union of faces of P that are orthogonal to τ . The set A_τ is a (possibly empty) smooth algebraic subvariety of M_P . It then follows from (95) and (101) that $H_P(\tau, \cdot)$ is constant on each connected component of A_τ . Indeed, on a component A^0 of A_τ , we have $H_P(\tau, \cdot) \equiv e^{-\langle \alpha^0, \tau \rangle}$, where α^0 is a lattice point in $\mu_P(A^0)$.

3.2.1. The normal bundle. Although equations (91) and (94) are necessary and sufficient conditions for $(0, w)$ to be a critical point of the phase on the τ -contour (as we shall show), they do not guarantee that the real part of the phase is maximal at that point. To obtain maximality, we need to strengthen condition (94) to require that $-\tau$ lies in the ‘normal bundle’ of P at w_z .

Recall that the normal cone C_F of a face F of P is given by

$$C_F = \{u \in \mathbb{R}^m : \langle u, x \rangle = \sup_{y \in P} \langle u, y \rangle, \quad \forall x \in F\}. \quad (102)$$

We now consider compact polytopes in $\mathbb{R}^m$ with arbitrary vertices that are not necessarily integral or even rational. We define the faces of such polytopes as in the integral case, and we use (102) to define the normal cones.

Definition: The normal bundle $\mathcal{N}(Q)$ of a (not necessarily integral) convex polytope $Q \subset \mathbb{R}^m$ is the subset of $T_{\mathbb{R}^m} = \mathbb{R}^m \times \mathbb{R}^m$ consisting of pairs (x, v) , where $x \in Q$ and v is in the normal cone C_F of the face F of Q that contains x . (Note that $\mathcal{N}(Q)$ is not a fiber bundle over Q .)

The normal bundle $\mathcal{N}(Q)$ is a piecewise smooth submanifold of $T_{\mathbb{R}^m}$; it is homeomorphic to $\mathbb{R}^m$ via the ‘exponential map’

$$\mathcal{E}_Q : \mathcal{N}(Q) \rightarrow \mathbb{R}^m, \quad \mathcal{E}_Q(x, v) = x + v, \quad x \in F, \quad v \in C_F \quad (F \text{ a face of } Q).$$

It is easily seen that $\mathcal{E}_Q$ is a homeomorphism and is a C^∞ (in fact, linear) diffeomorphism on each of the ‘pieces’ $\bar{F} \times C_F \subset \mathcal{N}(Q)$.

We shall use the following elementary, but not so well known, fact about the invertibility of Lipschitz maps. We let $B_\varepsilon(x_0) = \{x \in \mathbb{R}^m : |x - x_0| < \varepsilon\}$ denote the ε -ball about $x_0 \in \mathbb{R}^m$.

LEMMA 3.6. [Fan] *Let $f : U \rightarrow \mathbb{R}^m$ be a Lipschitz map, where U is open in $\mathbb{R}^m$. Then f has a local orientation-preserving Lipschitz inverse with Lipschitz constant L at a point $x_0 \in U$ if and only if there exists $\varepsilon > 0$ such that*

- i) $\liminf_{v \rightarrow 0} |f(x+v) - f(x)|/|v| \geq 1/L$ for all $x \in B_\varepsilon(x_0)$,
- ii) $\det f'(x) > 0$ for all $x \in B_\varepsilon(x_0)$ such that f is differentiable at x ,
- iii) $f(x_0) \notin f(\partial B_\varepsilon(x_0))$ and $\deg[f : \partial B_\varepsilon(x_0) \rightarrow \mathbb{R}^m \setminus \{f(x_0)\}] = 1$.

By the degree in (iii), we mean the degree of $f_*(\partial B_\varepsilon(x_0)) \in H_{m-1}(\mathbb{R}^m \setminus \{f(x_0)\}, \mathbb{Z})$. The hypotheses (i) and (iii) in [Fan] differ slightly from those above, but Fan's proof of sufficiency uses only (i)–(iii). (Necessity is obvious.) Hypothesis (i) is given as a lemma in [Fan]. Hypothesis (iii) above is replaced in [Fan] by the condition that the index of f at x_0 is 1, which is easily seen to be equivalent to (iii) under the assumptions (i) and (ii).

We now let

$$\mathcal{N}^\circ = \mathcal{N}(\tfrac{1}{p}P) \cap (\Sigma^\circ \times \mathbb{R}^m).$$

LEMMA 3.7. *For each point $z \in \mathbb{C}^{*m}$, there exists a unique $(x, v) \in \mathcal{N}^\circ$ such that*

$$\mu_\Sigma(e^{v/2} \cdot z) = x.$$

Proof. We note that the $\mathbb{R}_+^m$ action on $\mathbb{C}^{*m}$ descends via the moment map μ_Σ to an $\mathbb{R}_+^m$ action on Σ° , and we consider the map

$$\Phi : \mathcal{N}^\circ \rightarrow \Sigma^\circ, \quad \Phi(x, v) = \mu_\Sigma(e^{v/2} \cdot x).$$

It suffices to show that Φ is a bijection.

Let $\mathcal{L} : \Sigma^\circ \rightarrow \mathbb{R}^m$ be the diffeomorphism given by

$$\mathcal{L}(x) = \left(\log \frac{x_1}{1 - \sum x_j}, \dots, \log \frac{x_n}{1 - \sum x_j} \right), \quad (103)$$

so that

$$\mathcal{L} \circ \mu_\Sigma(z) = (\log |z_1|^2, \dots, \log |z_m|^2).$$

Thus, writing $x = \mu_\Sigma(z)$ we have

$$\begin{aligned} \mathcal{L} \circ \Phi(x, v) &= \mathcal{L} \circ \mu_\Sigma(e^{v/2} \cdot z) \\ &= (v_1 + \log |z_1|^2, \dots, v_m + \log |z_m|^2) \\ &= v + \mathcal{L}(x). \end{aligned} \quad (104)$$

We first observe that $\mathcal{L} \circ \Phi$ is proper: suppose on the contrary that the sequence $\{(x^n, v^n)\}$ is unbounded in $\mathcal{N}^\circ$, but $\mathcal{L} \circ \Phi(x^n, v^n) \rightarrow a \in \mathbb{R}^m$. By passing to a subsequence, we can assume that $x^n \rightarrow x^0 \in \frac{1}{p}P$. Then $x^0 \in \partial \Sigma$, since otherwise $v^n \rightarrow a - \mathcal{L}(x^0)$. Write $v^n = r^n u^n$, where $r^n > 0$, $|u^n| = 1$. We can assume without loss of generality that $u^n \rightarrow u^0$. We consider the case where $\Sigma x_j^0 < 1$, i.e., $\mu_\Sigma^{-1}(x^0)$ is not in the hyperplane at infinity. If $x^0 = 0$, then $u_j^0 \geq 0$ ($1 \leq j \leq m$), since otherwise $\mathcal{L} \circ \Phi(x^n, v^n)$ would diverge. But $u^0 \in C_{\{0\}}$; hence $\frac{1}{p}P \subset \{x : \langle u^0, x \rangle \leq 0\} \subset \mathbb{R}^m \setminus \Sigma^\circ$, a contradiction. Now suppose that $x_1^0 = \dots = x_k^0 = 0$, $x_l > 0$ for $k < l \leq m$. Then we conclude as before that $u_j^0 \geq 0$ for $1 \leq j \leq k$, and $u_l^0 = 0$ for $k < l \leq m$, and we again obtain a contradiction. Finally, if $\Sigma x_j^0 = 1$, we can change coordinates (permute the homogeneous coordinates in $\mathbb{CP}^m$) to reduce to the previous case.

Let $\mathcal{E} = \mathcal{E}_{\frac{1}{p}P} : \mathcal{N}(\frac{1}{p}P) \xrightarrow{\sim} \mathbb{R}^m$, and let $U = \mathcal{E}(\mathcal{N}^\circ) \subset \mathbb{R}^m$. We consider the map $f : U \rightarrow \mathbb{R}^m$ given by $f \circ \mathcal{E}|_{\mathcal{N}^\circ} = \mathcal{L} \circ \Phi$, i.e. by the commutative diagram:

$$\begin{array}{ccc} \mathcal{N}^\circ & \xrightarrow{\Phi} & \Sigma^\circ \\ \approx \downarrow \mathcal{E} & & \approx \downarrow \mathcal{L} \\ U & \xrightarrow{f} & \mathbb{R}^m \end{array}$$

Since $\mathcal{L} \circ \Phi$ is a proper map, f is also proper. Hence to show that Φ is a bijection, it suffices to show that f is a local homeomorphism and is therefore a (global) homeomorphism.

To describe the map f , for each $x \in \Sigma^\circ$, we let q_x denote the closest point in $\frac{1}{p}P$ to x . We note that $q_x \in \Sigma^\circ$; if $x \notin \frac{1}{p}P$, then $q_x \in \partial(\frac{1}{p}P)$; if $x \in \frac{1}{p}P$, then $q_x = x$. Furthermore, $\mathcal{E}^{-1}(x) = (q_x, x - q_x)$ and hence

$$f(x) = x - q_x + \mathcal{L}(q_x).$$

We shall show that f satisfies the hypotheses of Lemma 3.6. Let F be an arbitrary face of $\frac{1}{p}P \cap \Sigma^\circ$. To verify (i) and (ii), it suffices to show that $\det Df > 0$ on the (noncompact) polyhedron

$$U_F := \mathcal{E}(\bar{F} \times C_F) \cap U = \mathcal{E}[(\bar{F} \cap \Sigma^\circ) \times C_F].$$

To compute the determinant, we let $T_F \subset \mathbb{R}^m$, $N_F = T_F^\perp \subset \mathbb{R}^m$ denote the tangent and normal spaces, respectively, of F . Let $x^0 \in F$ be fixed. For $y \in U_F$, we have

$$y = x + v \xrightarrow{f} \mathcal{L}(x) + v, \quad x - x^0 \in T_F, \quad v \in N_F.$$

Choose orthonormal bases $\{Y_1, \dots, Y_r\}$, $\{Y_{r+1}, \dots, Y_m\}$ of T_F , N_F , respectively. We let Df denote the matrix of the derivative $(f|_{U_F})'$ with respect to the basis $\{Y_1, \dots, Y_m\}$ of $\mathbb{R}^m$. We have:

$$Df = \begin{pmatrix} T^t \mathcal{L}'(x) T & 0 \\ * & I \end{pmatrix}, \quad (105)$$

where T is the $m \times r$ matrix $[Y_1 \cdots Y_r]$. We have by (103),

$$(\mathcal{L}'(x))_{jk} = \frac{1}{x_0} + \delta_j^k \frac{1}{x_j}, \quad x_0 = 1 - \sum_{j=1}^m x_j > 0, \quad (106)$$

for $x \in \Sigma^\circ$. Hence $\mathcal{L}'(x)$ is a positive definite symmetric matrix, it being the sum of a semipositive matrix (all of whose entries are $\frac{1}{x_0}$) and a positive definite diagonal matrix. Therefore, $T^t \mathcal{L}'(x) T$ is positive definite, and hence $\det Df(x) = \det(T^t \mathcal{L}'(x) T) > 0$, completing the proof that hypotheses (i) and (ii) of Lemma 3.6 are satisfied.

Note that (106) implies that the eigenvalues of $\mathcal{L}'(x)$ are > 1 for $x \in \Sigma^\circ$ and hence by (105), $Df(x)$ is a diagonalizable matrix whose eigenvalues are real and ≥ 1 .

We verify (iii) by a homotopy argument: Choose a point $q^0 \in (\frac{1}{p}P)^\circ$. We contract $\frac{1}{p}P$ to q^0 ; i.e., for $0 \leq t \leq 1$, we let

$$Q_t = (1-t)\{q^0\} + \frac{t}{p}P,$$

so that $Q_0 = \{q^0\}$, $Q_1 = \frac{1}{p}P$. For $0 \leq t < 1$, $Q_t \subset \Sigma^\circ$, and hence we have a map

$$\Phi_t : \mathcal{N}(Q_t) \rightarrow \Sigma^\circ.$$

For $0 \leq t < 1$, we define $f_t : \mathbb{R}^m \rightarrow \mathbb{R}^m$ by the commutative diagram:

$$\begin{array}{ccc} \mathcal{N}(Q_t) & \xrightarrow{\Phi_t} & \Sigma^\circ \\ \approx \downarrow \mathcal{E}_{Q_t} & & \approx \downarrow \mathcal{L} \\ \mathbb{R}^m & \xrightarrow{f_t} & \mathbb{R}^m \end{array}$$

As before we have

$$f_t(x) = x - q_x^t + \mathcal{L}(q_x^t), \quad (107)$$

where q_x^t is the (unique) point of Q_t closest to x . The above argument shows that the maps f_t also satisfy (i) and (ii) of Lemma 3.6

We write

$$F : (\mathbb{R}^m \times [0, 1]) \cup (U \times \{1\}) \rightarrow \mathbb{R}^m, \quad (x, t) \mapsto f_t(x),$$

where $f_1 = f : U \rightarrow \mathbb{R}^m$. One easily sees that

$$|q_x^t - q_{x'}^{t'}| \leq |x - x'| + |t - t'|,$$

and hence F is continuous. Furthermore, F is uniformly continuous on $\mathbb{R}^m \times [0, t]$, for each $t < 1$.

Let H denote the set of $t \in [0, 1)$ such that $f_t : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is an orientation preserving homeomorphism. We note that $0 \in H$ since

$$f_0(x) = x - q + \mathcal{L}(q) = x + \text{constant}.$$

Note that (105) also holds for the maps f_t and hence as above, the eigenvalues of Df_t are real and ≥ 1 . Hence f_t satisfies conditions (i) and (ii) of Lemma 3.6 with $L = 1$, for all $t \in [0, 1]$.

First we show that H is open in $[0, 1)$. Suppose $t_0 \in H$. Then by Lemma 3.6, $\text{Lip}(f_{t_0}^{-1}) \leq 1$ and hence

$$|x - x_0| = 1 \implies |f_{t_0}(x) - f_{t_0}(x_0)| \geq 1.$$

By the uniform continuity of F on $\mathbb{R}^m \times [0, \frac{t_0+1}{2}]$, we can choose $\varepsilon > 0$ such that for all $x_0 \in \mathbb{R}^m$, we have

$$|f_t(x) - f_t(x_0)| \geq \frac{1}{2}, \quad \text{for } x \in \partial B_1(x_0), \quad |t - t_0| < \varepsilon. \quad (108)$$

To simplify notation, we shall write

$$\deg(f, x_0, r) := \deg[f : \partial B_r(x_0) \rightarrow \mathbb{R}^m \setminus \{f(x_0)\}].$$

We conclude from (108) that for $|t - t_0| < \varepsilon$,

$$\deg(f_t, x_0, 1) = \deg(f_{t_0}, x_0, 1) = 1,$$

and hence f_t is a local homeomorphism at x_0 , by Lemma 3.6. By the very first part of the argument that $f : U \rightarrow \mathbb{R}^m$ is proper, we easily see that f_t is proper. Since $x_0 \in \mathbb{R}^m$ is arbitrary, it follows that $f_t : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a covering map and therefore is a homeomorphism.

Next we show that H is closed in $[0, 1)$ and hence $H = [0, 1)$. Let $t_n \in H$ such that $t_n \rightarrow t_0 \in [0, 1)$. Since f_{t_0} satisfies condition (i) of Lemma 3.6, we can choose $\varepsilon > 0$ so that $f_{t_0}(x_0) \notin f_{t_0}(\partial B_\varepsilon(x_0))$. Then for n sufficiently large,

$$\deg(f_{t_0}, x_0, \varepsilon) = \deg(f_{t_n}, x_0, \varepsilon) = 1.$$

It follows as above that $f_{t_0} : \mathbb{R}^m \rightarrow \mathbb{R}^m$ is a homeomorphism.

We have shown that f_t is a homeomorphism for $0 \leq t < 1$. To complete the proof of the lemma, we must show that $f = f_1$ is a local homeomorphism. So we let $x_0 \in U$ be arbitrary, and we choose $\varepsilon > 0$ such that $\overline{B_\varepsilon(x_0)} \subset U$ and $f(x_0) \notin f(\partial B_\varepsilon(x_0))$. Then for $t < 1$ sufficiently close to 1, we have as before

$$\deg(f, x_0, \varepsilon) = \deg(f_t, x_0, \varepsilon) = 1.$$

Thus by Lemma 3.6, f is a local homeomorphism. □

As a consequence of Lemmas 3.5 and 3.7, we immediately obtain:

LEMMA 3.8. *For each $z \in \mathbb{C}^{*m}$, there exists $\tau_z \in \mathbb{R}^m$, $w_z \in M_P$ so that*

- $\mu_\Sigma(e^{\tau_z/2} \cdot z) = \frac{1}{p} \mu_P(w_z);$
- $(\frac{1}{p} \mu_P(w_z), -\tau_z) \in \mathcal{N}^\circ.$

Furthermore, τ_z, w_z are unique (modulo the $\mathbf{T}^m$ action on M_P) and also satisfy (91) and (94).

We shall show in §§3.2.2–3.2.3 that along the contour given by $\tau = \tau_z$, the phase $\Psi_\mathbb{C}$ has critical points and maximal real part at $(0, w_z)$.

We recall the concept of transition points mentioned in the introduction:

Definition: A point $z \in \mathbb{C}^{*m}$ is said to be a transition point if $-\tau_z$ is in the boundary of the normal cone of P at $\mu_P(w_z)$, or equivalently if z is in the boundary of one of the regions $\mathcal{R}_F$ described in the introduction.

Remark: It follows from (104) that $2 \text{Log}(\mathcal{R}_F) = \mathcal{L}(F) + C_F$, where $\text{Log}(z) = (\log |z_1|, \dots, \log |z_m|)$. Hence we can decompose $\mathbb{R}^m$ as the disjoint union of the sets $\mathcal{L}(F) + C_F$. If z is a transition point, then $2 \text{Log}(z)$ must lie in the common boundary of at least two of the sets $\mathcal{L}(F) + C_F$. Figure 4 below shows the transition points in log coordinates (as solid lines) for the case where P is the polytope of Figure 3.

Note that although $C_{F'}$ is orthogonal to F' in Figure 3, $C_{F'}$ is not orthogonal to $\mathcal{L}(F')$. Similarly, in Lemma 3.8, if $\mu_P(w_z) \in F'$, then the orbit $\mu_\Sigma(e^{r\tau_z} \cdot z)$ is not orthogonal to F' .




FIGURE 4. Transition points and regions in log coordinates

3.2.2. *Computation of the Hessian.* We consider

$$\mathcal{C}_z := \{e^{i\theta} \cdot w_z : e^{i\theta} \in \mathbf{T}^m\} = \{e^{i\theta} \cdot w_z : \theta \in T_F\} \subset M_P ,$$

which is a (totally real) submanifold of M_P of dimension r . We are going to show that $\{0\} \times \mathcal{C}_z \subset \mathbf{T}^m \times M_P$ is the set of critical points of $\Psi_{\mathbb{C}}$ with maximal real part along the τ_z contour. In order to apply the method of stationary phase as in the case where z is in the classically allowable region, one would also like to show that $\{0\} \times \mathcal{C}_z$ is a nondegenerate critical submanifold; i.e., the normal Hessian of the phase is nondegenerate along $\{0\} \times \mathcal{C}_z$. But the normal Hessian turns out to be degenerate whenever z is a transition point. For example, if $z \in \mu_{\Sigma}^{-1}(F)$, where F is a codimension r face of $\frac{1}{p}P$, then the corresponding normal Hessian has an r -dimensional nullspace (as our computations will show) and hence the normal Hessian is degenerate whenever $\mu_{\Sigma}(z) \in \partial(\frac{1}{p}P)$. However, we shall show:

LEMMA 3.9. *Let $z \in \mathbb{C}^{*m}$. Then $\{0\} \times \mathcal{C}_z$ is the (unique) component of the critical set of $\Psi_{\mathbb{C}}(\tau_z, \cdot, \cdot)$ where $\Re \Psi_{\mathbb{C}}$ attains its maximum on $\mathbf{T}^m \times M_P$. If z is not a transition point, then $\{0\} \times \mathcal{C}_z$ is a nondegenerate critical submanifold.*

Recall that in the previous section, we verified the proposition for the case where z is in the classically allowed region. So we now fix a point z in the classically forbidden region. Let F_z be the face of P containing $q_z := \mu_P(w_z)$, and let r denote the dimension of F_z . Note that $F - q_z$ an open subset of the tangent space T_F of $F \subset \mathbb{R}^m$.

Let $Y = \mu_P^{-1}(F_z)$, which is an r -dimensional complex submanifold, and thus has real dimension $2r$. We let $V \subset T_{M_P, w_z}$ be the tangent space to the $\mathbb{R}_+^m$ orbit of w_z . Recalling (98), we see that

$$V = \text{span}\{X_1(w_z), \dots, X_m(w_z)\} = \left\{ \sum a_j X_j(w_z) : (a_1, \dots, a_m) \in T_F \right\} ,$$

and hence $\dim V = r$. (Note that $V = JT^v$, where T^v is the tangent space to the fiber of μ_P .)

The normal bundle $\mathbf{N}$ to $\{0\} \times \mathcal{C}_z$ can be decomposed as follows:

$$\mathbf{N} = T_{\mathbf{T}^m, 0} \oplus V \oplus N_Y . \quad (109)$$

To prove Lemma 3.9, we need the following:

LEMMA 3.10. *The normal Hessian of $\Psi_{\mathbb{C}}(\tau_z, \cdot, \cdot)$ to $\{0\} \times \mathcal{C}_z$ with respect to a basis for $\mathbf{N}$ giving the decomposition (109) is of the form*

$$D_{\mathbf{N}}^2 \Psi_{\mathbb{C}}(\tau_z, 0, w_z) = \begin{pmatrix} A & B & 0 \\ B^t & 0 & 0 \\ 0 & 0 & C \end{pmatrix} ,$$

where

- i) A is an $m \times m$ negative-definite, real symmetric matrix;
- ii) B is an $m \times r$ pure imaginary matrix of rank r ;
- iii) C is a $(2m - 2r) \times (2m - 2r)$ real symmetric matrix, which is negative definite if z is not a transition point.

Proof. Let

$$D_{\mathbf{N}}^2 \Psi_{\mathbb{C}}(\tau_z, 0, w_z) = \begin{pmatrix} A & B & \star \\ B^t & \star'' & \star' \\ \star^t & \star'^t & C \end{pmatrix}.$$

To simplify notation, we now write $\tau = \tau_z$.

(i) To compute A , we see from (89) that

$$iD_{\varphi} \Psi_{\mathbb{C}}|_{\varphi=0} = \sum_{\alpha \in P} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2 \alpha + \text{a function of } z, \quad (110)$$

As in our previous computation, we then obtain

$$\begin{aligned} A &= \mu_P(e^{-\tau/2} \cdot w) \otimes \mu_P(e^{-\tau/2} \cdot w) - \sum_{\alpha \in P} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2 \alpha \otimes \alpha \\ &\quad + \mu_{p\Sigma}(e^{\tau/2} \cdot z) \otimes \mu_{p\Sigma}(e^{\tau/2} \cdot z) - \sum_{\alpha \leq p} |\widehat{m}_{\alpha}^{p\Sigma}(e^{\tau/2} \cdot z)|^2 \alpha \otimes \alpha. \end{aligned}$$

By Lemma 3.4, A is the sum of a negative semi-definite term and a strictly negative term, and hence is negative definite.

(ii) To compute B , we similarly obtain

$$iD_{\rho} D_{\varphi} \Psi_{\mathbb{C}}|_{\varphi=0} = \mu_P(e^{-\tau/2} \cdot w) \otimes \mu_P(e^{-\tau/2} \cdot w) - \sum_{\alpha \in P} |\widehat{m}_{\alpha}^P(e^{-\tau/2} \cdot w)|^2 \alpha \otimes \alpha, \quad (111)$$

where $D_{\rho} = (\partial/\partial\rho_1, \dots, \partial/\partial\rho_m) = (X_1, \dots, X_m)$. By Lemma 3.4,

$$(iD_{\rho} D_{\varphi} \Psi_{\mathbb{C}}|_{\varphi=0}, \lambda \otimes \lambda) > 0, \quad \text{for } \lambda \in T_F \setminus \{0\},$$

and hence B has rank r .

(iii) To determine C , we need to change variables. Let us recall that M_P is covered by affine coordinate charts centered at the vertices of P described as follows: Let $v = \alpha^0$ be a vertex of P . Since P is simplicial, there are exactly m edges (1-dimensional faces) incident to v . Choose $\alpha^1, \dots, \alpha^m$ on the respective edges. Let $\beta^j = \alpha^j - \alpha^0$; then $\{\beta^1, \dots, \beta^m\}$ form a basis for $\mathbb{R}^m$. Since we are assuming that P is Delzant, we can choose the α^j such that $\{\beta^1, \dots, \beta^m\}$ generate $\mathbb{Z}^m$. We consider the $m \times m$ matrix of integers

$$\Delta = (\Delta_{jk}) = (\beta_k^j).$$

Then $\det \Delta = \pm 1$ and the matrix $\Gamma := \Delta^{-1}$ has integer entries. Consider the (Zariski) neighborhood of v :

$$U_v := \{w \in M_P : \chi_v(w) \neq 0\} = \bigcup \{F : F \text{ is a face of } P, v \in \bar{F}\}.$$

(The U_v cover M_P .) Let $\eta_j = w^{\beta^j}$, $j = 1, \dots, m$. We note that the monomial mapping

$$\eta = (\eta_1, \dots, \eta_m) : \mathbb{C}^{*m} \rightarrow \mathbb{C}^{*m}$$

is bijective, its inverse given by

$$w_j = \eta_1^{\Gamma_{j1}} \dots \eta_m^{\Gamma_{jm}}.$$

Let $F^1, \dots, F^m$ denote the facets of P incident to v , indexed so that $\alpha_j \notin F^j$. Let $x \in F^j$ be arbitrary. We then have

$$\eta_j = \frac{\chi_{\alpha^j}(w)}{\chi_{\alpha^0}(w)} \rightarrow \frac{\chi_{\alpha^j}(x)}{\chi_{\alpha^0}(x)} = 0 \quad \text{as } \mu_P(w) \rightarrow x \in F^j.$$

(Since $\alpha^0 \in \bar{F}^j$, $\widehat{\chi}_{\alpha^0}(x) \neq 0$.) Therefore η extends to an isomorphism $\eta : (U_v, v) \xrightarrow{\sim} (\mathbb{C}^m, 0)$. Furthermore, the facets incident to v are the images of the divisors $\{\eta_j = 0\}$ under the moment map. (This shows that M_P is nonsingular if the Delzant condition is satisfied.)

We note that the standard unit vectors $e^j = \Gamma\beta^j = \Gamma\alpha^j - \Gamma v$. Consider the nonhomogeneous linear map

$$\tilde{\Gamma} : \mathbb{Z}^m \rightarrow \mathbb{Z}^m, \quad u \mapsto \Gamma u - \Gamma v.$$

Then $\tilde{\Gamma}(\alpha^0) = 0$, $\tilde{\Gamma}(\alpha^j) = e^j$ ($1 \leq j \leq m$). Consider the polytope $Q = \tilde{\Gamma}(P)$. Under the map $\tilde{\Gamma}$, the closed facets $\overline{F^j}$ correspond to polytopes in the coordinate hyperplanes $\{x_j = 0\}$. Hence the polytope Q lies in the upper quadrant $\{x_j \geq 0, 1 \leq j \leq m\}$. We also let

$$\tilde{\Delta} = \tilde{\Gamma}^{-1} : \mathbb{Z}^m \rightarrow \mathbb{Z}^m, \quad u \mapsto \Delta u + v.$$

Now let us rewrite the phase in terms of the η coordinates. We first note that

$$\chi_\alpha(w) = \chi_{\tilde{\Gamma}(\alpha)}(\eta), \quad w \in \mathbb{C}^{*m}, \quad \eta_j = w^{\beta^j}.$$

Hence,

$$H_P(\tau, w) = \frac{\sum_{\alpha \in P} e^{-\langle \alpha, \tau \rangle} |c_\alpha \chi_{\tilde{\Gamma}(\alpha)}(\eta)|^2}{\sum_{\alpha \in P} |c_\alpha \chi_{\tilde{\Gamma}(\alpha)}(\eta)|^2} = \frac{\sum_{\gamma \in Q} e^{-\langle \tilde{\Delta}(\gamma), \tau \rangle} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2}{\sum_{\gamma \in Q} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2}. \quad (112)$$

We first consider the case where $w_z = v$; i.e., $r = 0$. Expanding (112) in powers of η_j , we have

$$\begin{aligned} H_P(\tau, w) &= \frac{|c_v|^2 e^{-\langle v, \tau \rangle} + \sum_{j=1}^m |c_{\alpha^j}|^2 e^{-\langle \alpha^j, \tau \rangle} |\eta_j|^2 + \dots}{|c_v|^2 + \sum_{j=1}^m |c_{\alpha^j}|^2 |\eta_j|^2 + \dots} \\ &= e^{-\langle v, \tau \rangle} \left[1 + \sum_{j=1}^m c'_j \left(e^{-\langle \beta^j, \tau \rangle} - 1 \right) |\eta_j|^2 + \dots \right], \end{aligned}$$

where $c'_j = |c_{\alpha^j}/c_{\alpha^0}|^2 > 0$. Since $-\tau$ is in the normal cone at v , we have $\langle -\tau, v \rangle \geq \langle -\tau, \alpha_j \rangle$ and hence $\langle \beta_j, \tau \rangle \geq 0$, for $1 \leq j \leq m$. **Now suppose that z is not a transition point;** i.e., $-\tau$ is not in the boundary of the normal cone at $\{v\}$. Then τ is not perpendicular to any of the rays along the edges incident to v ; so $\langle \beta_j, \tau \rangle \neq 0$, and therefore $\langle \beta_j, \tau \rangle > 0$. Hence

$$\log H_P(\tau, w) = -\langle v, \tau \rangle + \sum_{j=1}^m b_j |\eta_j|^2 + \dots, \quad b_j = c'_j \left(e^{-\langle \beta^j, \tau \rangle} - 1 \right) < 0.$$

This leads to two conclusions: (i) $d \log H_P|_{w=w_z} = 0$ and hence $(0, w_z)$ is a critical point of $\Psi_{\mathbb{C}}(\tau, \cdot, \cdot)$; (ii) $C = D_w^2 \log H_P|_{w=w_z}$ is a diagonal matrix with negative eigenvalues $b_1, b_1, \dots, b_m, b_m$. Note that C is singular whenever z is in the boundary of the normal polyhedron of v .

Now consider the case $1 \leq r \leq m-1$. (If $r = m$, then z is in the allowable region.) We can assume without loss of generality that $Y = \mu_P^{-1}(F_z)$ is given by

$$\eta_1 = \dots = \eta_{m-r} = 0, \quad \eta_{m-r+1} \neq 0, \dots, \eta_m \neq 0. \quad (113)$$

In order to decompose the sum in (112), we consider the sets of lattice points

$$E_j := \{\gamma \in Q : \gamma_k = \delta_k^j \text{ for } 1 \leq k \leq m-r\} = Q \cap \{e^j + \sum_{l=m-r+1}^m s_l e^l\},$$

for $1 \leq j \leq m-r$. (These are the indices of the only terms that contribute to the normal Hessian at w_z .) We also let $E_0 = Q \cap (\{0_{m-r}\} \times \mathbb{Z}^r)$. Since $\langle \beta^l, \tau \rangle = 0$ for $l > m-r$, it follows that

$$\langle \tilde{\Delta}(\gamma), \tau \rangle = \langle \tilde{\Delta}(e^j), \tau \rangle = \langle \alpha^j, \tau \rangle \quad \forall \gamma \in E_j,$$

and similarly

$$\langle \tilde{\Delta}(\gamma), \tau \rangle = \langle \tilde{\Delta}(0), \tau \rangle = \langle v, \tau \rangle \quad \forall \gamma \in E_0.$$

We write

$$\eta = (\eta', \eta''), \quad \eta' = (\eta_1, \dots, \eta_{m-r}), \quad \eta'' = (\eta_{m-r+1}, \dots, \eta_m).$$

Let $E_j'' \subset \mathbb{Z}^m$, for $j = 0, 1, \dots, m-r$, be given by

$$\{0_{m-r}\} \times E_0'' = E_0, \quad \{0_{m-r}\} \times E_j'' = E_j - e^j \quad (1 \leq j \leq m-r).$$

Thus we rewrite (112):

$$\begin{aligned}
H_P(\tau, w) &= \frac{e^{-\langle v, \tau \rangle} \sum_{\gamma \in E_0} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2 + \sum_{j=1}^{m-r} \left[e^{-\langle \alpha^j, \tau \rangle} \sum_{\gamma \in E_j} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2 \right] + \dots}{\sum_{\gamma \in E_0} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2 + \sum_{j=1}^{m-r} \left[\sum_{\gamma \in E_j} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2 \right] + \dots} \\
&= e^{-\langle v, \tau \rangle} \frac{\sum_{\nu \in E_0''} c'_{0\nu} |\chi_\nu(\eta'')|^2 + \sum_{j=1}^{m-r} \left[\sum_{\nu \in E_j''} c'_{j\nu} |\chi_\nu(\eta'')|^2 \right] e^{-\langle \beta^j, \tau \rangle} |\eta_j|^2 + \dots}{\sum_{\nu \in E_0''} c'_{0\nu} |\chi_\nu(\eta'')|^2 + \sum_{j=1}^{m-r} \left[\sum_{\nu \in E_j''} c'_{j\nu} |\chi_\nu(\eta'')|^2 \right] |\eta_j|^2 + \dots} \\
&= e^{-\langle v, \tau \rangle} \left[1 + \sum_{j=1}^{m-r} \frac{\lambda_j(\eta'')}{\lambda_0(\eta'')} \left(e^{-\langle \beta^j, \tau \rangle} - 1 \right) |\eta_j|^2 + O(\|\eta'\|^4) \right], \tag{114}
\end{aligned}$$

where $c'_{j\nu} \in \mathbb{R}_+$ and

$$\lambda_j(\eta'') = \sum_{\nu \in E_j''} c'_{j\nu} |\chi_\nu(\eta'')|^2 = c'_{j0} + \sum_{\nu \in E_j'' \setminus \{0\}} c'_{j\nu} |\chi_\nu(\eta'')|^2 > 0 \quad (0 \leq j \leq m-r).$$

(Note that (114) implies that $H_P(\tau, w) = e^{-\langle v, \tau \rangle}$ for $w \in F_z$, as we observed earlier, and that $(0, w_z)$ is a critical point of $\Psi_{\mathbb{C}}$.) **Suppose as before that z is not a transition point;** i.e., $-\tau$ is not in the boundary of the normal cone at F_z . Then $\langle \beta_j, \tau \rangle \neq 0$ and in fact $\langle \beta_j, \tau \rangle > 0$, for $1 \leq j \leq m-r$, since $-\tau$ is in the cone at F_z . It now follows from (114) that $C = D_{\eta'}^2 \log H_P|_{\{\eta'=0\}}$ is diagonal (with respect to the η' coordinates) with negative eigenvalues.

(iv) To complete the proof of the lemma, we now show that the other blocks of the normal Hessian $D_{\mathbf{N}}^2 \Psi_{\mathbb{C}}$ vanish. We first note that since $H_P(\tau_z, \cdot)$ is constant on Y , the matrix $\star''$ vanishes. We also immediately conclude from (114) that

$$\star' = D_{\rho''} D_{\eta'} \log H_P|_{\{\eta'=0\}} = 0,$$

where $\rho'' = (\Re \eta_{m-r+1}, \dots, \Re \eta_m)$.

It remains to show that $\star = 0$. Assuming (113), we expand (110) as before:

$$\begin{aligned}
iD_{\varphi} \Psi_{\mathbb{C}}|_{\varphi=0} &= \frac{\sum_{\gamma \in Q} e^{-\langle \tilde{\Delta}(\gamma), \tau \rangle} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2 \tilde{\Delta}(\gamma)}{\sum_{\gamma \in Q} |c_{\tilde{\Delta}(\gamma)} \chi_\gamma(\eta)|^2} + f(z) \\
&= e^{-\langle v, \tau \rangle} \left[\tilde{g}_0(\eta'') + \sum_{j=1}^{m-r} |\eta_j|^2 \tilde{g}_j(\eta'') + O(\|\eta'\|^4) \right] + f(z).
\end{aligned}$$

Hence

$$\star = D_{\eta'} D_{\varphi} \Psi_{\mathbb{C}}|_{\varphi=0, \eta'=0} = 0.$$

□

3.2.3. Proof of Lemma 3.9. It follows from (114) that $(0, w_z)$ is a critical point of $\Psi_{\mathbb{C}}(\tau, \cdot, \cdot)$ and hence $\{0\} \times \mathcal{C}_z$ is contained in the critical set. We conclude from Lemma 3.10 that the real $r \times r$ matrix $B^t A^{-1} B = (iB)^t (-A^{-1})(iB)$ is positive definite, and hence

$$\det D_{\mathbf{N}}^2 \Psi_{\mathbb{C}} = \det A \det(B^t A^{-1} B) \det C \neq 0, \tag{115}$$

if z is not a transition point. Therefore $\{0\} \times \mathcal{C}_z$ is a nondegenerate critical manifold of $\Psi_{\mathbb{C}}$ whenever z is not a transition point.

Recalling (97), we set

$$b(z) := -\Psi_{\mathbb{C}}(\tau_z, 0, w_z) = -p \log H_{\Sigma}(-\tau_z, z) - \log H_P(\tau_z, w_z). \tag{116}$$

We now show that the maximum of $\Re \Psi_{\mathbb{C}}$ on the τ -contour is attained on $\{0\} \times \mathcal{C}_z$ and hence the maximum equals $-b(z)$. We recall that by (87), it suffices to show that $\Re \Psi(\tau, 0, w)$ attains its maximum at $w = w_z$, and by (97), this is equivalent to establishing that $\log H(\tau, \cdot)$, whose Hessian is C , takes its maximum at w_z .

We first suppose that z is not a transition point. Note that by the above, $H(\tau, \cdot)$ is constant and has a local maximum along $Y = \mu_P^{-1}(F_z)$. Suppose on the contrary that there is a point $\tilde{w} \in M_P$ such that

$$H(\tau, \tilde{w}) = \sup_{w \in M_P} H(\tau, w) > H(\tau, w_z). \quad (117)$$

Let $\tilde{F}$ denote the face that contains $\mu_P(\tilde{w})$. Since $H(\tau, \cdot)$ is constant on F_z , by (117) the face $\tilde{F}$ is not contained in $\overline{F_z}$, and hence $C_{\tilde{F}} \cap C_{F_z}^\circ = \emptyset$, where C_F° denotes the nonboundary points of the cone C_F . Since $-\tau \in C_{F_z}^\circ$ (by the assumption that z is not a transition point), it follows that $-\tau \notin C_{\tilde{F}}$. Then we conclude from (114) that at least one of the eigenvalues of $C(\tilde{w})$ is positive, contradicting (117). On the other hand, if z is a transition point, then $-\tau \in C_{F'}^\circ$, where F' is a face whose closure contains F_z . Then $H(\tau, w_z) = H(\tau, w')$ for $w' \in F'$ and $H(\tau, \cdot)$ attains a local maximum along F' . Then by the above argument, $H(\tau, w')$ is a global maximum. Recalling (116), we then have

$$-b(z) = \Psi_{\mathbb{C}}(\tau_z, 0, w_z) = \sup_{w \in M_P} \Psi_{\mathbb{C}}(\tau_z, 0, w) = \sup_{\varphi \in \mathbf{T}^m, w \in M_P} \Re \Psi_{\mathbb{C}}(\tau_z, \varphi, w_z). \quad (118)$$

This completes the proof of Lemma 3.9. $\square$

3.2.4. Asymptotic expansions on the classically forbidden region. We continue the proof of Proposition 3.1(ii). Equation (118) says that $\Psi + b(z)$ is a phase function of positive type that takes the value 0 on the critical set $\{0\} \times \mathcal{C}_z$. **If z is not a transition point**, the points of $\{0\} \times \mathcal{C}_z$ are nondegenerate critical points. As before, we obtain by the method of stationary phase an asymptotic expansion

$$K(z) = N^{\frac{m+r}{2}} e^{-Nb(z)} [c_0^F(z) + c_1^F(z)N^{-1} + c_2^F(z)N^{-2} + \dots] \quad (119)$$

valid over each open region $\mathcal{R}_F^\circ$. (Since the critical submanifold $\{0\} \times \mathcal{C}_z$ has codimension $3m - r$, the leading term of the expansion of $e^{Nb(z)}K(z)$ contains $N^{2m - \frac{1}{2}(3m-r)} = N^{\frac{m+r}{2}}$.) To show that $\Pi_{|NP}(z, z)$ also has the asymptotic expansion (119), we must show that

$$S_N(z) = O(N^{-k} e^{-Nb(z)}) \quad \forall k \in \mathbb{N}, \quad (120)$$

where the bound is locally uniform in $\mathcal{C}^j$ for all j . We deform the $\mathbf{T}^m$ integral in (71) to an integral over $\{|\zeta| = \tau_z\}$ as before, so that we have

$$\begin{aligned} S_N(z) &= \frac{1}{(2\pi)^m} \int_{M_P} \int_{M_P} \int_{\mathbf{T}^m} \tilde{R}_N(w, \eta) \left(\sum_{|\alpha| \leq p} e^{\langle \alpha, \tau_z + i\varphi \rangle} |\tilde{m}_\alpha^{p\Sigma}(z)|^2 \right)^N \\ &\quad \times \left(\sum_{\alpha \in P} e^{-\langle \alpha, \tau_z + i\varphi \rangle} \hat{m}_\alpha^P(\eta) \overline{\hat{m}_\alpha^P(w)} \right)^N d\varphi dw d\eta, \end{aligned} \quad (121)$$

where $\tilde{R}_N = \frac{(Np+m)!}{(Np)!} R_N$ is rapidly decaying. We first note that

$$\left| \sum_{|\alpha| \leq p} e^{\langle \alpha, \tau_z + i\varphi \rangle} |\hat{m}_\alpha^{p\Sigma}(z)|^2 \right| \leq \sum_{|\alpha| \leq p} e^{\langle \alpha, \tau_z \rangle} |\hat{m}_\alpha^{p\Sigma}(z)|^2 = H_{p\Sigma}(-\tau_z, z).$$

By the Cauchy-Schwartz inequality and the fact established above that $H_P(\tau_z, w)$ takes its maximum at $w = w_z$, we have

$$\begin{aligned} \left| \sum_{\alpha \in P} e^{-\langle \alpha, \tau_z + i\varphi \rangle} \hat{m}_\alpha^P(\eta) \overline{\hat{m}_\alpha^P(w)} \right| &\leq \sum_{\alpha \in P} e^{-\langle \alpha, \tau_z/2 \rangle} |\hat{m}_\alpha^P(\eta)| e^{-\langle \alpha, \tau_z/2 \rangle} |\hat{m}_\alpha^P(w)| \\ &\leq \left(\sum_{\alpha \in P} e^{-\langle \alpha, \tau_z \rangle} |\hat{m}_\alpha^P(\eta)|^2 \right)^{\frac{1}{2}} \left(\sum_{\alpha \in P} e^{-\langle \alpha, \tau_z \rangle} |\hat{m}_\alpha^P(w)|^2 \right)^{\frac{1}{2}} \\ &= H_P(\tau_z, \eta)^{\frac{1}{2}} H_P(\tau_z, w)^{\frac{1}{2}} \leq H_P(\tau_z, w_z). \end{aligned}$$

Therefore, we can write

$$S_N(z) = \frac{1}{(2\pi)^m} \int_{M_P} \int_{M_P} \int_{\mathbf{T}^m} \tilde{R}_N(w, \eta) e^{N\hat{\Psi}(\varphi, w, \eta; z)} d\varphi dw d\eta ,$$

where the phase satisfies

$$\Re \hat{\Psi} \leq \log [H_{p\Sigma}(-\tau_z, z) H_P(\tau_z, w_z)] = e^{-b(z)} .$$

The estimate (120) now follows from the fact that $\tilde{R}_N$ is rapidly decaying.

Thus, we have verified the asymptotic formula (ii) of Proposition 3.1, together with (a) and (b), and we have shown that $b(z)$ is given by (116).

Remark: It follows from (72), (87), and (118) that in a neighborhood $U \subset \subset \mathbb{C}^{*m}$ of a transition point z^0 , we have uniform bounds of the form

$$\Pi_{|NP}(z, z) \leq C_U N^{2m} e^{-Nb(z)} . \quad (122)$$

But a sharper inequality should hold; for example, we know that if $z^0 \in \partial \mathcal{A}_P$, then $b(z^0) = 0$ and

$$\Pi_{|NP}(z^0, z^0) \leq \Pi_{N_P}^{\mathbb{CP}^m}(z^0, z^0) = [p^m + o(1)] N^m .$$

It is an open question whether we have asymptotic expansions of the form (119) at transition points. See §5.3 for a discussion of the issues involved in studying the asymptotics at transition points.

3.2.5. The decay function. In this section we verify (c)–(e) of Proposition 3.1(ii) and then prove Proposition 3.2. We first note that when $z \in \mathcal{A}_P$, we have $\tau_z = z$ and hence by (116), $b = 0$ on $\mathcal{A}_P$. Furthermore, since $z \mapsto \tau_z$ is easily seen to be continuous on $\mathbb{C}^{*m}$ and $\mathcal{C}^\infty$ on each $\overline{\mathcal{R}_F}$, the same holds for $b(z)$.

We now show that b is $\mathcal{C}^1$ by computing its derivative. Recalling that $b(z) = -\Psi_{\mathbb{C}}(\tau_z, 0, w_z; z)$, we have

$$-D_z b = D_\tau \Psi_{\mathbb{C}}|_{(\tau_z, 0, w_z, z)} \cdot D_z \tau_z + D_w \Psi_{\mathbb{C}}|_{(\tau_z, 0, w_z, z)} \cdot D_z w_z + D_z \Psi_{\mathbb{C}}|_{(\tau_z, 0, w_z, z)} .$$

Here, $D_z \tau_z$ and $D_z w_z$ are only piecewise continuous, being discontinuous at transition points. By (88)–(91) and the fact that $(0, w_z)$ is a critical point of $\Psi_{\mathbb{C}}(\tau_z, \cdot, \cdot)$, we have

$$D_\tau \Psi_{\mathbb{C}}|_{(\tau_z, 0, w_z, z)} = -i D_\varphi \Psi_{\mathbb{C}}|_{(\tau_z, 0, w_z, z)} = 0 .$$

The critical point condition also says that $D_w \Psi_{\mathbb{C}}|_{(\tau_z, 0, w_z, z)} = 0$. Recalling (97), we thus have

$$D_z b = -D_z \Psi_{\mathbb{C}}(\tau, 0, w; z)|_{\tau=\tau_z, w=w_z} = -p D_z \log H_\Sigma(-\tau, z)|_{\tau=\tau_z} \in \mathcal{C}^0(\mathbb{C}^{*m}) . \quad (123)$$

Therefore, $b \in \mathcal{C}^1(\mathbb{C}^{*m})$, verifying (e).

To verify (c), we need to find a more explicit formula for the derivative, using the log coordinates $\zeta_j = \rho_j + i\theta_j = \log z_j$. We consider the map $\xi = (\xi_1, \dots, \xi_m) : \mathbb{C}^{*m} \rightarrow \mathbb{C}^{*m}$ given by

$$\xi(z) = e^{\tau_z/2} \cdot z = e^{\tau_z/2 + \rho + i\theta} , \quad (124)$$

so that $\mu_\Sigma \circ \xi(z) = \frac{1}{p} \mu_P(w_z)$. By (123),

$$\begin{aligned} D_\rho b &= -p D_\rho \log H_\Sigma(-\tau, e^\rho)|_{\tau=\tau_z} \\ &= -p D_\rho \log \left(1 + \sum e^{\tau_j + 2\rho_j} \right) \Big|_{\tau=\tau_z} + p D_\rho \log \left(1 + \sum e^{2\rho_j} \right) . \end{aligned} \quad (125)$$

We have

$$\begin{aligned} D_\rho \log \left(1 + \sum e^{\tau_j + 2\rho_j} \right) \Big|_{\tau=\tau_z} &= \left(\frac{2e^{\tau_1 + 2\rho_1}}{1 + \sum e^{\tau_j + 2\rho_j}}, \dots, \frac{2e^{\tau_m + 2\rho_m}}{1 + \sum e^{\tau_j + 2\rho_j}} \right) \\ &= \left(\frac{2|\xi_1|^2}{1 + \|\xi\|^2}, \dots, \frac{2|\xi_m|^2}{1 + \|\xi\|^2} \right) = 2\mu_\Sigma \circ \xi , \end{aligned} \quad (126)$$

and similarly, $D_\rho \log (1 + \sum e^{2\rho_j}) = 2\mu_\Sigma$. Hence

$$D_\rho b = 2p(\mu_\Sigma - \mu_\Sigma \circ \xi) . \quad (127)$$

Let $z \in \mathbb{C}^{*m} \setminus \overline{\mathcal{A}_P}$. If z is not a transition point, then the asymptotic expansion holds at z , and as we observed at the beginning of §3.2, $\Pi_{|NP}(z, z)$ is rapidly decaying; hence $b(z) > 0$. Now suppose on the

contrary that z is a transition point and $b(z) = 0$. Then z would be a critical point for b , which is impossible since $Db|_z \neq 0$ whenever $z \notin \overline{\mathcal{A}_P}$ by (127). This completes the proof of (c).

It remains to verify formula (10). First we establish (11): Fix a point z in the classically forbidden region, and make the change of variables $\rho_j = \sigma_j/2 + \log|z_j|$. Then by (127),

$$b(e^{\tau_z/2} \cdot z) - b(z) = \int_0^{\tau_z} D_\sigma b(e^{\sigma/2} \cdot z) \cdot d\sigma = p \int_0^{\tau_z} [\mu_\Sigma(e^{\sigma/2} \cdot z) - \mu_\Sigma \circ \xi(e^{\sigma/2} \cdot z)] \cdot d\sigma. \quad (128)$$

By definition, $q(z) = \mu_P(w_z) = p\mu_\Sigma \circ \xi(z)$, and hence $p\mu_\Sigma \circ \xi(e^{\sigma/2} \cdot z) = q(e^{\sigma/2} \cdot z)$. Furthermore $b(e^{\tau_z/2} \cdot z) = 0$ since $e^{\tau_z/2} \cdot z = \xi(z) \in \partial\mathcal{A}_P$, and formula (11) follows from (128).

Note that the integral in (11) is independent of the path from 0 to τ_z . Choosing the path $\sigma(r) = r\tau_z$, $0 \leq r \leq 1$, we have

$$\int_0^{\tau_z} q(e^{\sigma/2} \cdot z) \cdot d\sigma = \int_0^{\tau_z} q(z) \cdot d\sigma = \langle q(z), \tau_z \rangle. \quad (129)$$

On the other hand, since $D_\sigma \log(1 + \|e^{\sigma/2} \cdot z\|^2) = \mu_\Sigma(e^{\sigma/2} \cdot z)$, we have

$$\int_0^{\tau_z} \mu_\Sigma(e^{\sigma/2} \cdot z) \cdot d\sigma = \log(1 + \|e^{\tau_z/2} \cdot z\|^2) - \log(1 + \|z\|^2). \quad (130)$$

Substituting (129)–(130) in (11), we obtain formula (10). This completes the proof of part (ii) of Proposition 3.1.

We note that b fails to be $\mathcal{C}^2$ at transition points. This follows from (127), (124), and the fact that $z \mapsto \tau_z$ is piecewise $\mathcal{C}^1$ with discontinuous derivative at the interfaces between the regions $\mathcal{R}_F$.

Remark: An alternate derivation of (10) is as follows: By an argument similar to the above, we obtain

$$\log H_P(\tau, w) = \int_0^\tau \mu_P(e^{\sigma/2} \cdot w) \cdot d\sigma, \quad (131)$$

and therefore

$$\log H_P(\tau_z, w_z) = \langle -\mu_P(w_z), \tau_z \rangle. \quad (132)$$

Formula (10) is an immediate consequence of (116) and (132).

We now prove Proposition 3.2: By Proposition 3.1(ii) and (116),

$$\frac{1}{N} \log \Pi_{|NP}(z, z) \rightarrow -b(z) = p \log H_\Sigma(-\tau_z, z) + \log H_P(\tau_z, w_z), \quad (133)$$

for all non-transition points z . Let

$$u_N(z) = \frac{1}{N} \log \Pi_{|NP}(z, z) + p \log(1 + \|z\|^2) = \frac{1}{N} \log \sum_{\alpha \in NP} \binom{Np}{\alpha} e^{2\langle \alpha, \rho \rangle}, \quad z \in \mathbb{C}^{*m}. \quad (134)$$

Thus for all non-transition points z , we have

$$u_N(z) \rightarrow u_\infty(z) := p \log(1 + \|z\|^2) - b(z) = p \log(1 + \|e^{\tau_z/2} \cdot z\|^2) + \log H_P(\tau_z, w_z), \quad (135)$$

We must show that convergence of (135) also holds at the transition points and is uniform on compact subsets of $\mathbb{C}^{*m}$. We again use the log coordinates $\zeta_j = \rho_j + i\theta_j = \log z_j$, so that

$$D_\rho u_N = \frac{1}{N} D_\rho \log \sum_{\alpha \in NP} \binom{Np}{\alpha} e^{2\langle \alpha, \rho \rangle} = \frac{\sum_{\alpha \in NP} \binom{Np}{\alpha} e^{2\langle \alpha, \rho \rangle} 2\alpha}{N \sum_{\alpha \in NP} \binom{Np}{\alpha} e^{2\langle \alpha, \rho \rangle}} = \frac{2}{N} \mu_{NP}, \quad (136)$$

and therefore $\|D_\rho u_N\| \leq 2p$. Since $\{u_N(z^0)\}$ converges for any non-transition point z^0 , it follows that $\{u_N\}$ is uniformly bounded and uniformly equicontinuous on compact sets. Therefore it converges uniformly on compact sets in $\mathbb{C}^{*m}$. $\square$

3.2.6. *Precise asymptotics on the classically allowed region.* In this section, we prove part (i) of Proposition 3.1. Since

$$\Pi_{Np\Sigma}(z, z) \equiv \frac{\dim H^0(\mathbb{CP}^m, \mathcal{O}(Np))}{\text{Vol}(\mathbb{CP}^m)} = \binom{Np+m}{m} m! = \prod_{j=1}^m (Np+j),$$

we have by (29) and (39),

$$\Pi_{|NP}(z, z) = \prod_{j=1}^m (Np+j) + R_N(z),$$

where

$$R_N(z) = - \sum_{\alpha \in Np\Sigma \setminus NP} \frac{1}{\|\chi_\alpha\|^2} |\hat{\chi}_\alpha(z)|^2 = - \left[\prod_{j=1}^m (Np+j) \right] \sum_{\alpha \in Np\Sigma \setminus NP} |\hat{m}_\alpha^{Np\Sigma}(z)|^2. \quad (137)$$

Since $\#(Np\Sigma \setminus NP) = O(N^m)$, Proposition 3.1(i) is an immediate consequence of (137) and the following uniform asymptotic decay estimate for monomials.

LEMMA 3.11. *Suppose that $K \subset \mathbb{C}^{*m}$ is compact, and let $U \subset \Sigma$ be an open neighborhood of $\mu_\Sigma(K)$. Then there exist positive constants $C_l = C_{l,K,U}$, $\lambda = \lambda_{K,U}$ such that*

$$\|\hat{m}_\alpha^{N\Sigma}\|_{C^l(K)} \leq C_l e^{-\lambda N} \quad \text{whenever } \frac{\alpha}{N} \in \Sigma \setminus U, \quad N \geq 1, l \geq 0.$$

Proof. We first prove the $\mathcal{C}^0$ estimate. Choose a positive integer $k > 2m^2/\text{dist}(\Sigma \setminus U, \mu_\Sigma(K))$. We consider the finite collection of polytopes

$$Q_\beta := 2m\Sigma + \beta \subset k\Sigma, \quad \beta \in \mathbb{Z}^m \cap (k-2m)\Sigma.$$

For each β as above, we let $b_\beta(z)$ be the function on $\mathbb{C}^{*m}$ given by formula (10) with P replaced by Q_β and with p replaced by k . We claim that the $\mathcal{C}^0$ estimate holds with

$$\lambda = \frac{1}{k+\varepsilon} \inf \left\{ b_\beta(z) : z \in K, \beta \in \mathbb{Z}^m \cap (k-2m)\Sigma, \frac{1}{k}Q_\beta \subset \Sigma \setminus U \right\} > 0.$$

Suppose on the contrary that the estimate does not hold with this value of λ . Then we can choose an increasing sequence of integers $\{N_j\}$ and a sequence $\alpha^j \in N(\Sigma \setminus U) \cap \mathbb{Z}^m$ such that

$$e^{\lambda N_j} \sup_K \left| \hat{m}_{\alpha^j}^{N_j \Sigma} \right|^2 \rightarrow +\infty. \quad (138)$$

By passing to a subsequence, we can assume that $\frac{\alpha^j}{N_j} \rightarrow x \in \Sigma \setminus U$. Choose $\beta \in \mathbb{Z}^m \cap (k-2m)\Sigma$ such that

$$x \in \text{interior}_\Sigma \left(\frac{1}{k}Q_\beta \right),$$

and hence $k\alpha^j \in N_j Q_\beta$ for $j \gg 1$. Since $\text{dist}(x, K) \geq \frac{2m^2}{k} \geq \text{diam}(\frac{1}{k}Q_\beta)$, it follows that $(\frac{1}{k}Q_\beta) \cap K = \emptyset$. Hence by Proposition 3.2 applied to the polytope Q_β , there is a positive constant A such that

$$\Pi_{|NQ_\beta}(z, z) \leq A \exp \left(-\frac{k}{k+\varepsilon} b_\beta(z) N \right) \leq A e^{-\lambda k N} \quad \forall z \in K, \forall N \geq 1. \quad (139)$$

We shall use the multinomial inequality:

$$\binom{N}{\alpha}^k \leq \binom{kN}{k\alpha}. \quad (140)$$

One way to verify (140) is to regard $\binom{N}{\alpha}$ as the partition function $\mathcal{P}_N(\alpha)$ for the polytope Σ ; i.e., the number of ways to write $\alpha = \beta^1 + \dots + \beta^N$, where $\beta \in \Sigma$ (see §2.1). Now, given $k\alpha \in kN\Sigma$ we first write $k\alpha = \alpha + \dots + \alpha$ (k times) and then decompose each α as above, resulting in $\binom{N}{\alpha}^k$ different decompositions of $k\alpha$ as a sum of kN elements of Σ . This gives us the lower bound for $\binom{kN}{k\alpha}$ in (140).

We thus have

$$|\widehat{m}_{\alpha^j}^{N_j \Sigma}|^{2k} = \left(\frac{N_j}{\alpha^j}\right)^k |\widehat{\chi}_{\alpha^j}|^{2k} \leq \left(\frac{kN_j}{k\alpha^j}\right) |\widehat{\chi}_{k\alpha^j}|^2 = |\widehat{m}_{k\alpha^j}^{kN_j \Sigma}|^2.$$

Since $k\alpha^j \in N_j Q_\beta$, it then follows from (139) that

$$|\widehat{m}_{\alpha^j}^{N_j \Sigma}(z)|^2 \leq |\widehat{m}_{k\alpha^j}^{kN_j \Sigma}|^{2/k} \leq \Pi_{|N_j Q_\beta}(z, z)^{1/k} \leq A^{1/k} e^{-\lambda N_j},$$

for all $z \in K$, which contradicts (138). Hence the $\mathcal{C}^0$ estimate holds.

We now verify the $\mathcal{C}^1$ estimate: We have

$$|\widehat{m}_\alpha(z)| = \left(\frac{N}{\alpha}\right)^{\frac{1}{2}} h(z)^{\frac{N}{2}} |z^\alpha|, \quad h(z) = \frac{1}{1 + \|z\|^2}.$$

Differentiating, we obtain

$$D_\rho |\widehat{m}_\alpha(z)| = \left(\frac{N}{\alpha}\right)^{\frac{1}{2}} h(z)^{\frac{N}{2}} |z^\alpha| \left[\alpha + \frac{N}{2} D_\rho \log h \right] = |\widehat{m}_\alpha(z)| \left[\alpha + \frac{N}{2} D_\rho \log h \right],$$

and hence the $\mathcal{C}^1$ estimate follows from the $\mathcal{C}^0$ estimate. Differentiating repeatedly, we obtain all the $\mathcal{C}^l$ estimates. $\square$

The proof of Proposition 3.1 is now complete.

4. DISTRIBUTION OF ZEROS

In order to deduce Theorem 3 from Proposition 3.1, we need to relate $\mathbf{E}_{|NP}(Z_s)$ to the Szegő kernel $\Pi_{|NP}$. The idea is essentially the same as in [SZ1, Prop. 3.1], but neither the statement nor the proof there cover the application we need. Hence we provide a more general statement for the zeros of general linear systems on compact Kähler manifolds (Proposition 4.1).

4.1. Expected distribution of zeros and Szegő kernels. As in §1.2, we let $L \rightarrow M$ be a holomorphic line bundle over a compact Kähler manifold, and we consider a linear subspace $\mathcal{S} \subset H^0(M, L^N)$, which we endow with a Hermitian inner product. (The set of zero divisors $\{Z_s : s \in \mathcal{S}\} \equiv \mathbb{P}\mathcal{S}$ is called a *linear system* on M .) We let e_L be a local holomorphic frame over a trivializing chart U . We choose an orthonormal basis $\{S_j : 1 \leq j \leq k\}$ for $\mathcal{S}$ and we write $S_j = f_j e_L$ over U . Any section $s \in \mathcal{S}$ may then be written as

$$s = \langle c, F \rangle e_L^{\otimes N}, \quad \text{where } F = (f_1, \dots, f_k), \quad \langle c, F \rangle = \sum_{j=1}^k c_j f_j.$$

The Kodaira map is given by:

$$\Phi_{\mathcal{S}} : M \dashrightarrow \mathbb{CP}^{k-1}, \quad \Phi_{\mathcal{S}}(z) = [f_1(z), \dots, f_k(z)]. \quad (141)$$

(We use the symbol ‘ $\dashrightarrow$ ’ in (141) to indicate that the map $\Phi_{\mathcal{S}}$ is a rational map; it is holomorphic if and only if $\mathcal{S}$ has no basepoints other than fixed components.) Under a change of orthonormal basis, one gets a unitarily equivalent map. The current of integration over the zeros of $s = \langle c, F \rangle e_L^{\otimes N}$ is then given locally by the Poincaré-Lelong formula:

$$Z_s = \frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \log |\langle c, F \rangle|. \quad (142)$$

It is of course independent of the choice of local frame e_L and basis $\{S_j\}$.

We write a section locally as $s = f e_L$ and we write its Hermitian norm as $\|s(z)\|_h = a(z)^{-\frac{1}{2}} |f(z)|$ where $a(z) = \|e_L(z)\|_h^{-2}$. The *curvature form* $c_1(L, h)$ of L is given locally by

$$c_1(L, h) = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log a.$$

PROPOSITION 4.1. *Let (L, h) be a Hermitian line bundle on a compact complex manifold M . Let $\mathcal{S}$ be a subspace of $H^0(M, L)$ of dimension ≥ 2 . We give $\mathcal{S}$ an inner product and we let γ be the associated Gaussian probability measure. Then the expected zero current of a random section $s \in \mathcal{S}$ is given by*

$$\begin{aligned} \mathbf{E}_\gamma(Z_s) &= \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \Pi_{\mathcal{S}}(z, z) + c_1(L, h) \\ &= \Phi_{\mathcal{S}}^* \omega_{\text{FS}} + D_0, \end{aligned}$$

where D_0 is the fixed component of the linear system $\mathbb{P}\mathcal{S}$.

Proof. The Gaussian measure γ is the standard Gaussian for an inner product on $\mathcal{S}$. Let $\{S_j\}$ be an orthonormal basis for this inner product. As above, we choose a local nonvanishing section e_L of L over $U \subset M$, and we write

$$s = \sum_{j=1}^k c_j S_j = \langle c, F \rangle e_L,$$

where $S_j = f_j e_L$, $F = (f_1, \dots, f_k)$. As in the proof of [SZ1], Proposition 3.1, we then write $F(x) = |F(x)|u(x)$ so that $|u| \equiv 1$ and

$$\log |\langle c, F \rangle| = \log |F| + \log |\langle c, u \rangle|. \quad (143)$$

By (142), we have

$$\begin{aligned} (\mathbf{E}_\gamma(Z_s), \varphi) &= \frac{\sqrt{-1}}{\pi} \int_{\mathbb{C}^k} (\partial \bar{\partial} \log |\langle c, F \rangle|, \varphi) d\gamma(c) \\ &= \frac{\sqrt{-1}}{\pi} \int_{\mathbb{C}^k} (\partial \bar{\partial} \log |F|, \varphi) d\gamma(c) + \frac{\sqrt{-1}}{\pi} \left(\int_{\mathbb{C}^k} \partial \bar{\partial} \log |\langle c, u \rangle| d\gamma(c), \varphi \right), \end{aligned}$$

for all test forms $\varphi \in \mathcal{D}^{m-1, m-1}(U)$. Upon integration in c , the second term of (143) becomes constant in z and the derivatives kill it. The first term is independent of c so we may remove the Gaussian integral. Thus

$$\mathbf{E}_\gamma(Z_s) = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log |F|^2 = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \left(\log \sum \|S_j\|^2 + \log a \right). \quad (144)$$

Recalling that $\Pi_{\mathcal{S}}(z, z) = \sum \|S_j(z)\|^2$ and that $c_1(L, h) = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log a$, the first identity of the proposition follows.

Before completing the proof of the second identity, we recall the definition of the pull-back of a form by a rational map $X \xrightarrow{g} Y$: we let $\tilde{\Gamma}$ be the desingularization of the graph $\Gamma \subset X \times Y$ of g so that we have $g \circ \pi_1 = \pi_2$, where π_1, π_2 are the projections from $\tilde{\Gamma}$ to X, Y respectively. Then for any smooth form η on Y we define the pull-back current $g^* \eta := \pi_{1*} \pi_2^* \eta$, which is easily seen to have $\mathcal{L}^1$ coefficients with singularities on the indeterminacy locus I_g of g .

We can suppose that U is chosen so that there is a $\psi \in \mathcal{O}(U)$ with $\text{Div}(\psi) = D_0$ on U . Write $f_j = \psi g_j$. Then by (144) we have

$$\mathbf{E}_\gamma(Z_s) = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log |F|^2 = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log |\psi|^2 + \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \sum |g_j|^2.$$

The first term equals D_0 . By definition, the second term equals $\Phi_{\mathcal{S}}^* \omega_{\text{FS}}$ outside the indeterminacy locus $I_{\Phi_{\mathcal{S}}}$. Since $I_{\Phi_{\mathcal{S}}}$ has real codimension ≥ 4 and $\log \sum |g_j|^2$ is (pluri-)subharmonic, the coefficients of $\frac{i}{2\pi} \partial \bar{\partial} \log \sum |g_j|^2$ are measures that do not give mass to $I_{\Phi_{\mathcal{S}}}$; i.e., the coefficients are in $\mathcal{L}^1(U)$. Since the same is true for the coefficients of $\Phi_{\mathcal{S}}^* \omega_{\text{FS}}$, the current equality extends to all of U . $\square$

We note that the expected zero current $\mathbf{E}_\gamma(Z_s)$ is a smooth form outside the base point set of $\mathcal{S}$. We shall use the following result on simultaneous expected zeros (away from base points).

PROPOSITION 4.2. *Let M be a compact complex manifold, and let $(L_1, h_1), \dots, (L_k, h_k)$ be Hermitian line bundles on M ($1 \leq k \leq \dim M$). Suppose we are given subspaces $\mathcal{S}_j \subset H^0(M, L_j)$ with inner products $\langle \cdot, \cdot \rangle_j$ and let γ_j denote the associated Gaussian probability measures on the $\mathcal{S}_j$ ($1 \leq j \leq k$). Let U be an open*

subset of M on which $\mathcal{S}_j$ has no base points for all j . Then the expected simultaneous-zero current of random sections $s_1 \in \mathcal{S}_1, \dots, s_k \in \mathcal{S}_k$ is given over U by

$$\mathbf{E}_{\gamma_1 \times \dots \times \gamma_k}(Z_{s_1, \dots, s_k}) = \bigwedge_{j=1}^k \mathbf{E}_{\gamma_j}(Z_{s_j}) = \bigwedge_{j=1}^k \left[\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \Pi_{\mathcal{S}_j}(z, z) + c_1(L_j, h_j) \right] = \bigwedge_{j=1}^k \Phi_{\mathcal{S}_j}^* \omega_{\text{FS}}.$$

Proof. We first note that the expect current is well defined, since for all choices of the s_j the total mass of the zero current $Z_{s_1, \dots, s_k}$ equals the Chern number $c_1(L_1) \cdots c_1(L_k)$, and hence for each test form $\varphi \in \mathcal{D}^{m-k, m-k}(U)$, the function $(Z_{s_1, \dots, s_k}, \varphi)$ is in $\mathcal{L}^\infty(\mathcal{S}_1 \times \dots \times \mathcal{S}_k)$.

Next we note the following functorial property of the expected zero current: Suppose that $\rho : Y \rightarrow M$ is a holomorphic mapping of complex manifolds and let $\rho^* \gamma_j$ be the pulled-back Gaussian measure on $\rho^* \mathcal{S}_j \subset H^0(Y, \rho^* L)$. Then

$$\mathbf{E}_{\rho^* \gamma_j}(Z_{t_j}) = \rho^* \mathbf{E}_{\gamma_j}(Z_{s_j}) \quad \text{on } \rho^{-1}(U) \quad (t_j \in \rho^* \mathcal{S}_j). \quad (145)$$

Indeed, (145) follows immediately from Proposition 4.1 and its proof. As a special case, suppose that Y is a submanifold of M and ρ is the inclusion. Then (145) becomes

$$\mathbf{E}_{\gamma_j}([Y] \wedge Z_{s_j}) = [Y] \wedge \mathbf{E}_{\gamma_j}(Z_{s_j}) \quad \text{on } U. \quad (146)$$

(In fact, (146) holds for any subvariety Y , but we don't need this fact here.)

We now verify the current identity by induction on k . For $k = 1$, this is Proposition 4.1, so assume that $k > 1$ and the proposition has been verified for $k - 1$. Consider $Y = |Z_{s_1, \dots, s_{k-1}}|$. By Bertini's Theorem, we know that Y is smooth (and of codimension $k - 1$) for almost all $s_1, \dots, s_{k-1}$. Therefore by (146),

$$\int Z_{s_1, \dots, s_k} d\gamma_k(s_k) = \int Z_{s_1, \dots, s_{k-1}} \wedge Z_{s_k} d\gamma_k(s_k) = Z_{s_1, \dots, s_{k-1}} \wedge \mathbf{E}_{\gamma_k}(Z_{s_k}) \quad \text{for a.a. } s_1, \dots, s_{k-1}.$$

Integrating over $s_1, \dots, s_{k-1}$ and using the inductive hypothesis, we obtain

$$\mathbf{E}_{\gamma_1 \times \dots \times \gamma_k}(Z_{s_1, \dots, s_k}) = \mathbf{E}_{\gamma_1 \times \dots \times \gamma_{k-1}}(Z_{s_1, \dots, s_{k-1}}) \wedge \mathbf{E}_{\gamma_k}(Z_{s_k}) = \mathbf{E}_{\gamma_1}(Z_{s_1}) \wedge \dots \wedge \mathbf{E}_{\gamma_k}(Z_{s_k}).$$

The other equalities follow from Proposition 4.1. $\square$

If we consider the case where $M = \mathbb{CP}^m$, we obtain a formula for the expected simultaneous zero current for random polynomials with given Newton polytopes:

COROLLARY 4.3. *The expected zero current of k independent random polynomials $f_j \in H^0(\mathbb{CP}^m, \mathcal{O}(p_j), P_j)$, $1 \leq j \leq k$, is given over $\mathbb{C}^{*m}$ by*

$$\mathbf{E}_{|P_1, \dots, P_k|}(Z_{f_1, \dots, f_k}) = \bigwedge_{j=1}^k \left(\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \Pi_{|P_j|}(z, z) + p_j \omega_{\text{FS}}^{\mathbb{CP}^m} \right) \quad \text{on } \mathbb{C}^{*m}.$$

Proof. Since $H^0(\mathbb{CP}^m, \mathcal{O}(p_j), P_j)$ has no base points in $\mathbb{C}^{*m}$, the conclusion follows from Proposition 4.2 and the fact that $c_1(\mathcal{O}(p_j), h_{\text{FS}}) = p_j \omega_{\text{FS}}$. $\square$

Remark: By Proposition 4.1, the identity of currents

$$\mathbf{E}_{|P|}(Z_f) = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log \Pi_{|P|}(z, z) + p \omega_{\text{FS}}^{\mathbb{CP}^m} \quad (147)$$

holds on all of $\mathbb{CP}^m$.

We say that p is *minimal* for P if

$$\min_{\alpha \in P} \alpha_j = 0 \quad (1 \leq j \leq n), \quad \max_{\alpha \in P} |\alpha| = p. \quad (148)$$

(Otherwise, we can choose p to be smaller, after translating the polytope if necessary.) The hypothesis that p is minimal for P guarantees that $H^0(\mathbb{CP}^m, \mathcal{O}(p), P)$ has no fixed divisors. In this case, (147) holds as an identity of $(1, 1)$ -forms with $\mathcal{L}_{\text{loc}}^1$ coefficients.

4.2. Asymptotics of zeros. In this section, we prove Theorems 3 and 4 and give some additional asymptotic formulas for zeros. Theorem 1 is a special case of Corollaries 5 and 6 of Theorem 4.

To prove Theorem 3, we let u_N, u_∞ be as in the proof of Proposition 3.2. By Corollary 4.3 (recalling that $\omega_{\text{FS}} = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} \log(1 + \|z\|^2)$), we have

$$\frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} u_N = \frac{1}{N} \mathbf{E}_{|NP}(Z_f) \quad \text{on } \mathbb{C}^{*m}, \quad (149)$$

and hence u_N is plurisubharmonic. By (135) and (149), we conclude that

$$\frac{1}{N} \mathbf{E}_{|NP}(Z_f) \rightarrow \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} u_\infty = p\omega_{\text{FS}} - \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} b, \quad (150)$$

where differentiation is in the distribution sense.

We now show that the current $\partial\bar{\partial} b \in \mathcal{D}'^{1,1}(\mathbb{C}^{*m})$ is given by a $(1,1)$ -form with piecewise smooth coefficients; in fact $\partial\bar{\partial} b$ is $\mathcal{C}^\infty$ on each of the regions $\mathcal{R}_F$ given by (9). (Equivalently, the Radon measure $\partial\bar{\partial} b \wedge \omega_{\text{FS}}$ does not charge the set of transition points, and hence formula (150) can be interpreted as differentiation in the ordinary sense on the regions $\mathcal{R}_F$.) By Proposition 3.1, $b \in \mathcal{C}^1(\mathbb{C}^{*m})$; i.e., if $z^0 \in \partial\mathcal{R}_F \cap \partial\mathcal{R}_{F'}$, then the values of $db(z^0)$ computed in the two regions $\mathcal{R}_F$ and $\mathcal{R}_{F'}$ agree. Then for a test form $\varphi \in \mathcal{D}^{m-1, m-1}(\mathbb{C}^{*m})$, we have

$$(\partial\bar{\partial} b, \varphi) = \sum_F \int_{\mathcal{R}_F} b \partial\bar{\partial} \varphi = \sum_F \int_{\mathcal{R}_F^\circ} \partial\bar{\partial} b \wedge \varphi - \sum_F \int_{\partial\mathcal{R}_F} (\bar{\partial} b \wedge \varphi + b \wedge \partial\varphi). \quad (151)$$

Note that $\partial\mathcal{R}_F$ consists of $\mathcal{C}^\infty$ real hypersurfaces (consisting of those points of $\partial\mathcal{R}_F$ that are contained in the boundary of only one other region $\mathcal{R}_{F'}$) together with submanifolds of real codimension ≥ 2 , and hence Stokes' Theorem applies (see e.g. [Fe, 4.2.14]). Since b and $\bar{\partial} b$ are continuous on $\mathbb{C}^{*m}$, the boundary integral terms in (151) cancel out and we obtain

$$(\partial\bar{\partial} b, \varphi) = \sum_F \int_{\mathcal{R}_F^\circ} \partial\bar{\partial} b \wedge \varphi = \int_{\mathbb{C}^{*m} \setminus E} \partial\bar{\partial} b \wedge \varphi, \quad (152)$$

where $E := \bigcup_F \partial\mathcal{R}_F$ is the set of transition points. Formula (152) says that the current $\partial\bar{\partial} b$ is a $(1,1)$ -form with piecewise smooth coefficients obtained by differentiating b on the regions $\mathcal{R}_F^\circ$.

We now let

$$\psi_P = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} u_\infty = p\omega_{\text{FS}} - \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} b \quad (153)$$

on each of the regions $\mathcal{R}_F$. By (152), the current ψ_P is also a piecewise smooth $(1,1)$ -form; by (150)

$$N^{-1} \mathbf{E}_{|NP}(Z_f) \rightarrow \psi_P \quad (\text{weakly}). \quad (154)$$

We shall show $\mathcal{L}_{\text{loc}}^1$ convergence of (154) when we prove Theorem 4 below. Continuing with the proof of Theorem 3, we observe that (ii) is an immediate consequence of (153), since $b = 0$ on the classically allowed region.

To verify (iii), we again use the log coordinates $\rho_j + i\theta_j = \log z_j$, so that

$$u_\infty = p \log \left(1 + \sum e^{2\rho_j} \right) - b(e^\rho). \quad (155)$$

Since u_∞ depends only on $(\rho_1, \dots, \rho_m)$, we have

$$\psi_P = \frac{\sqrt{-1}}{2\pi} \partial\bar{\partial} u_\infty = \frac{\sqrt{-1}}{8\pi} \sum_{jk} \frac{\partial^2 u_\infty}{\partial \rho_j \partial \rho_k} d\zeta_j \wedge d\bar{\zeta}_k \geq 0. \quad (156)$$

Thus we must show that the Hessian of $u_\infty(\rho)$ has rank r at points $z^0 \in \mathcal{R}_F^\circ$, where $r = \dim F$. From (125)–(126) and (155), we obtain

$$D_\rho u_\infty = 2p \mu_\Sigma \circ \xi, \quad (157)$$

where $\xi(z) = e^{\tau_z/2} \cdot z$ as before. Hence $\frac{1}{2p} D_\rho^2 u_\infty$ equals the Jacobian of $\mu_\Sigma \circ \xi$. Since $\mu_\Sigma \circ \xi(z) = \frac{1}{p} \mu_P(w_z)$, we easily see that

$$\mu_\Sigma \circ \xi : \mathcal{R}_F^\circ \rightarrow F, \quad (158)$$

so $\text{rank } D(\mu_\Sigma \circ \xi) \leq r$. In fact, (158) is a submersion, so that the rank equals r . To see that it is a submersion, we recall from the proof of Lemma 3.7 that the ‘lipeomorphism’ $\Phi^{-1} : \Sigma^\circ \rightarrow \mathcal{N}^\circ$ restricts to a diffeomorphism

$$\Sigma^\circ \supset \mu_\Sigma(\mathcal{R}_F^\circ) \xrightarrow{\sim} F \times C_F \subset \mathcal{N}^\circ, \quad \mu_\Sigma(z) \mapsto \left(\frac{1}{p} \mu_P(w_z), -\tau_z \right) = (\mu_\Sigma \circ \xi(z), -\tau_z),$$

and hence (158) is a submersion, completing the proof of (iii).

We now prove Theorem 4; the case $k = 1$ of the theorem will then yield part (i) of Theorem 3.

Let $P_1, \dots, P_k$ be Delzant polytopes, as in Theorem 4. We first show that

$$N^{-k} \mathbf{E}_{|NP}(Z_{f_1, \dots, f_k}) \rightarrow \psi_k := \psi_{P_1} \wedge \dots \wedge \psi_{P_k} \quad (\text{weakly}). \quad (159)$$

For $1 \leq j \leq k$, we let

$$u_N^j = \frac{1}{N} \log \Pi_{|NP_j}(z, z) + p_j \log(1 + \|z\|^2),$$

so that, recalling (135),

$$u_N^j(z) \rightarrow u_\infty^j(z) := p_j \log(1 + \|z\|^2) - b(z) = p_j \log(1 + \|e^{\tau_z^j/2} \cdot z\|^2) + \log H_{P_j}(\tau_z^j, w_z^j),$$

where τ_z^j, w_z^j are as in Lemma 3.8 with $P = P_j$. By (153),

$$\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_\infty^j = \psi_{P_j}.$$

Recalling Corollary 4.3, we introduce the (k, k) -forms

$$\kappa_N := \left(\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_N^1 \right) \wedge \dots \wedge \left(\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_N^k \right) = N^{-k} \mathbf{E}_{|NP_1, \dots, NP_k}(Z_{f_1, \dots, f_k}). \quad (160)$$

Since $u_N^j \rightarrow u_\infty^j$ locally uniformly, it follows from the Bedford-Taylor Theorem [BT, Kl] that

$$\kappa_N \rightarrow \left(\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_\infty^1 \right) \wedge \dots \wedge \left(\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_\infty^k \right) \quad (\text{weakly}). \quad (161)$$

In fact, the limit current in (161) is absolutely continuous, and hence is equal to the locally bounded (piecewise smooth) (k, k) -form ψ_k . To see this, we note that by the Bedford-Taylor Theorem,

$$\bigwedge_{j=1}^k (\psi_{P_j} * \varphi_\epsilon) = \bigwedge_{j=1}^k \left[\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} (u_\infty^j * \varphi_\epsilon) \right] \rightarrow \bigwedge_{j=1}^k \left(\frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_\infty^j \right),$$

where φ_ϵ denotes an approximate identity. Since the currents ψ_{P_j} have coefficients in $\mathcal{L}_{\text{loc}}^\infty$, the forms $\bigwedge_{j=1}^k (\psi_{P_j} * \varphi_\epsilon)$ have locally uniformly bounded coefficients; absolute continuity of the limit current follows. The weak limit (159) now follows from (160)–(161).

To complete the proof of Theorem 4, we must show that $\kappa_N \rightarrow \psi_k$ in $\mathcal{L}^1(K)$ for all compact $K \subset \mathbb{C}^{*m}$. Let $\varepsilon > 0$ be arbitrary, and choose a nonnegative function $\eta \in \mathcal{D}(\mathbb{C}^{*m})$ such that $\eta \equiv 1$ on a neighborhood of $(E_1 \cup \dots \cup E_k) \cap K$, where E_j is the set of transition points for P_j , and

$$\int_{\mathbb{C}^{*m}} \psi_k \wedge \eta \omega_{\text{FS}}^{m-k} < \varepsilon.$$

By (159)–(160),

$$\int_{\mathbb{C}^{*m}} \kappa_N \wedge \eta \omega_{\text{FS}}^{m-k} < 2\varepsilon \quad \text{for } N \gg 0.$$

(Note that the above integrands are nonnegative.) Therefore,

$$\|\kappa_N - \psi_k\|_{\mathcal{L}^1(K)} \leq \|(1 - \eta)(\kappa_N - \psi_k)\|_{\mathcal{L}^1(K)} + \|\eta \kappa_N\|_{\mathcal{L}^1(K)} + \|\eta \psi_k\|_{\mathcal{L}^1(K)}.$$

Since κ_N is a positive (k, k) -form,

$$\|\eta\kappa_N\|_{\mathcal{L}^1(K)} = \int_K \kappa_N \wedge \eta\omega_{\text{FS}}^{m-k} < 2\varepsilon.$$

Similarly, $\|\eta\psi_k\|_{\mathcal{L}^1(K)} = \int_K \psi_k \wedge \eta\omega_{\text{FS}}^{m-k} < \varepsilon$. Since

$$(1 - \eta)(\kappa_N - \psi_k) \rightarrow 0 \quad \text{uniformly on } K,$$

it follows that

$$\|\kappa_N - \psi_k\|_{\mathcal{L}^1(K)} \rightarrow 0,$$

completing the proof of Theorems 3 and 4. $\square$

We note that the conclusion of Theorem 4 can be replaced with an asymptotic expansion away from transition points:

THEOREM 4.4. *Let $P_1, \dots, P_k$ be Delzant polytopes. Let U be a relatively compact domain in $\mathbb{C}^{*m}$ such that $\overline{U}$ does not contain transition points for any of the P_j . Then we have a complete asymptotic expansion of the form*

$$\frac{1}{N^k} \mathbf{E}_{|NP_1, \dots, NP_k}(Z_{f_1, \dots, f_k}) \sim \psi_{P_1} \wedge \dots \wedge \psi_{P_k} + \frac{\varphi_1}{N} + \dots + \frac{\varphi_n}{N^n} + \dots \quad \text{on } U,$$

with uniform C^∞ remainder estimates, where the φ_j are smooth (k, k) -forms on U .

Proof. By (134) and (160) we have

$$\frac{1}{N^k} \mathbf{E}_{|NP_1, \dots, NP_k}(Z_{f_1, \dots, f_k}) = \bigwedge_{j=1}^k \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \left[\frac{1}{N} \log \Pi_{|NP_j}(z, z) + p_j \log(1 + \|z\|^2) \right].$$

The conclusion now follows from the asymptotic expansion of Proposition 3.1. $\square$

The proof of Theorem 3(iii) gives us some more information about the expected zero current in the classically forbidden region, which we state in the theorem below. We first define the *complexified normal cones*

$$\tilde{C}_F = \{\tau + i\theta : \tau \in C_F, \theta \in T_F^\perp\}.$$

(Recall that $C_F \subset T_F^\perp$.) We note that $\tilde{C}_F$ is a semi-group, which acts on $\mathcal{R}_F$ by the rule $\eta(z) = e^\eta \cdot z$; we call this action the ‘(joint) normal flow.’ The (maximal) orbits of the normal flow are of the form $\tilde{C}_F \cdot z^0 = \{e^\eta \cdot z^0 : \eta \in \tilde{C}_F\}$, where $z^0 \in \mu_\Sigma^{-1}(F)$. We note that the orbit $\tilde{C}_F \cdot z^0$ is a complex $(m - r)$ -dimensional submanifold (with boundary) of $\mathbb{C}^{*m}$. (Indeed, $(\tilde{C}_F \cdot z^0) \cap \mathcal{R}_F^\circ$ is a submanifold without boundary in $\mathcal{R}_F^\circ$.)

THEOREM 4.5. *Let P be a Delzant polytope and let ψ_P be the limit expected zero current of Theorem 3. Then ψ_P vanishes along the orbits of the normal flow.*

Proof. Let

$$O := \tilde{C}_F \cdot z^0 = \{e^{\tau+i\theta} \cdot z^0 : \tau \in C_F, \theta \in T_F^\perp\}$$

be a maximal orbit of the normal flow, where $\mu_\Sigma(z^0) \in F$. For $z = e^{\tau+i\theta} \cdot z^0 \in O$, we have

$$\mu_\Sigma \circ \xi(z) = \frac{1}{p} \mu_P(w_z) = \frac{1}{p} \mu_P(w_{e^\tau \cdot z^0}) = \frac{1}{p} \mu_P(w_{z^0}) = \mu_\Sigma(z^0)$$

and hence $\mu_\Sigma \circ \xi$ is constant on O . It then follows from (156)–(157) that $\psi_P|_O = 0$. $\square$

Remark: Theorem 4.5 tells us about the behavior of tangent directions to typical zero sets $|Z_f|$ of polynomials $f \in H^0(\mathbb{CP}^m, \mathcal{O}(Np), NP)$ as $N \rightarrow \infty$. Roughly speaking, the tangent spaces of $|Z_f|$ are highly likely to be close to containing the tangent spaces of the orbits of the normal flow, for large N . E.g., let us consider a region $\mathcal{R}_F^\circ$, where F is an edge (i.e., $\dim F = 1$). Let T_z^{flow} denote the holomorphic tangent space to

the normal flow through a point $z \in \mathcal{R}_F^\circ$. Then $\dim_{\mathbb{C}} T_z^{\text{flow}} = \dim_{\mathbb{C}} Z_f = m - 1$, and for any compact set $K \subset \mathcal{R}_F^\circ$, we have

$$\frac{1}{N} \mathbf{E}_{|NP} \left(\int_{Z_f \cap K} \text{dist}(T_{Z_f, z}^{1,0}, T_z^{\text{flow}})^2 d\text{Vol}_{Z_f}(z) \right) \longrightarrow 0, \quad (162)$$

where ‘dist’ means the distance in the projective space of complex $(m-1)$ -dimensional subspaces of $T_{\mathbb{C}P^m, z}^{1,0}$.

To verify (162), we let V be a compactly supported vector field on $\mathcal{R}_F^\circ$ such that for all $z \in \mathcal{R}_F^\circ$, V_z is a $(1,0)$ -vector tangent to the normal flow through z . Let $\varphi = *(iV \wedge \bar{V}) \in \mathcal{D}^{m-1, m-1}(\mathcal{R}_F^\circ)$ be given by

$$\lambda \wedge \varphi_z = (\lambda, iV_z \wedge \bar{V}_z) \frac{1}{m!} \omega_{\text{FS}}^m, \quad \lambda \in T_z^{*,1} \mathbb{C}^m, \quad z \in \mathcal{R}_F^\circ.$$

By Theorem 4.5, we have

$$(\psi_P, \varphi) = \int \psi_P \wedge \varphi = \int (\psi_P, iV_z \wedge \bar{V}_z) \frac{1}{m!} \omega_{\text{FS}}^m = 0.$$

We let $\mathcal{T}_f \in \mathcal{C}^\infty(Z_f^{\text{reg}}, T_{Z_f}^{m-1, m-1})$ denote the dual to the volume form on Z_f^{reg} , so that

$$\int_{Z_f} \|\mathcal{T}_f \wedge V \wedge \bar{V}\| d\text{Vol}_{Z_f} = \int_{Z_f} (\mathcal{T}_f, \varphi) d\text{Vol}_{Z_f} = (Z_f, \varphi). \quad (163)$$

Hence by Theorem 3(i),

$$\frac{1}{NP} \mathbf{E}_{|NP} \left(\int_{Z_f} \|\mathcal{T}_f \wedge V \wedge \bar{V}\| d\text{Vol}_{Z_f} \right) = \frac{1}{N} \mathbf{E}_{|NP} (Z_f, \varphi) \rightarrow (\psi_P, \varphi) = 0, \quad (164)$$

which yields (162).

As mentioned in the introduction, we can apply Proposition 4.2 (as in [SZ1] for the case $k = 1$) to the asymptotic expansion of the Szegő kernel in [Ze] to immediately obtain the following asymptotics using toric norms:

PROPOSITION 4.6. *Let (M_P, L_P) be as above, and let I_j, θ_j be the action-angle variables of the moment map $\mu_P : M_P \rightarrow \mathbb{R}^m$ of the $\mathbf{T}^m$ -action on M_P . Then we have:*

$$\frac{1}{N^k} E_{\gamma_{N^k}^{M_P}}(Z_{f_1, \dots, f_k}) = \omega_P^k + O\left(\frac{1}{N}\right) = \left(\sum_{j=1}^m dI_j \wedge d\theta_j \right)^k + O\left(\frac{1}{N}\right).$$

Proof. Let $\omega_P = \sum_{j=1}^m dI_j \wedge d\theta_j$ be the natural Kähler form on M_P . We recall the asymptotic formula for the Szegő kernel in [Ze]:

$$\frac{1}{N} \Phi_{L_P^N}^* \omega_{\text{FS}} = \omega_P + O(1/N).$$

By Proposition 4.2 with $M = M_P$, $L = L_P^N$, we then have

$$\mathbf{E}_{M_P}(Z_{s_1, \dots, s_k}) = (\mathbf{E}_{M_P}(Z_s))^k = \Phi_{L_P^N}^* \omega_{\text{FS}}^k = N^k [\omega_P^k + O(1/N)].$$

□

4.3. Amoebas in the plane. The term ‘amoeba’ was introduced by Gelfand, Kapranov and Zelevinsky [GKZ] to refer to the image under the moment map of a zero set $Z_{f_1, \dots, f_k}$ of polynomials, and have been studied in various contexts (see [FPT, GKZ, Mi1, PR] and the references in the survey article by Mikhalkin [Mi2]). The image of a zero set under the moment map μ_Σ is called a *compact amoeba*, while the image under the map

$$\text{Log} : \mathbb{C}^m \rightarrow \mathbb{R}^m, \quad (z_1, \dots, z_m) \mapsto (\log |z_1|, \dots, \log |z_m|)$$

is a *noncompact amoeba*, or simply an *amoeba*. Note that Log is the moment map for the $\mathbf{T}^m$ action with respect to the Euclidean symplectic form $\sum dx_j \wedge dy_j$, and $\text{Log} = \frac{1}{2} \mathcal{L} \circ \mu_\Sigma$, where $\mathcal{L} : \Sigma^\circ \approx \mathbb{R}^m$ is the diffeomorphism given by (103).

To illustrate what our statistical results can say about amoebas, we consider zero sets in $\mathbb{C}^{*2}$. An amoeba in $\mathbb{R}^2$ is the image of a plane algebraic curve under the Euclidean moment map Log . An example of an

amoeba of the form $\text{Log}(Z_f)$, where f is a quartic polynomial in two variables with (full) Newton polytope 4Σ , is given in the illustration from [Th] reproduced in Figure 5 below.²




FIGURE 5. An amoeba with polytope 4Σ

One notices that this amoeba contains 12 ‘tentacles’. By definition, a tentacle on a compact amoeba A is a connected component of a small neighborhood in A of $A \cap \partial\Sigma$; the tentacles of a noncompact amoeba correspond to those of the compact amoeba under the diffeomorphism $\Sigma^\circ \approx \mathbb{R}^m$.

There is a natural injective map from the set of connected components (which are convex sets) of the complement of a noncompact amoeba A to the set $P \cap \mathbb{Z}^2$ of lattice points of the polytope, and that there are amoebas for which each lattice point is assigned to a component of the complement. (This fact is also valid in higher dimensions; see [FPT, Mi1].) For a generic 2-dimensional noncompact amoeba with polytope P , each lattice point in ∂P corresponds to a distinct unbounded component of $\mathbb{R}^2 \setminus A$, and adjacent lattice points correspond to adjacent unbounded components. (The correspondence is given by [Mi1, §3.1] or [FPT].) Hence the number of tentacles of A equals the number of points of $\partial P \cap \mathbb{Z}$; this number is called the *length* of ∂P . (For example, the 12 tentacles in Figure 5 correspond to the 12 lattice points of $\partial(4\Sigma)$.) Each tentacle corresponds to a segment connecting 2 adjacent lattice points on ∂P .

We can decompose ∂P into two pieces: $\partial^\circ P = \partial P \cap p\Sigma^\circ$ and $\partial^e P = P \cap \partial(p\Sigma)$. Tentacles corresponding to segments of $\partial^\circ P$ end (in the compact picture Σ) at a vertex of Σ , and tentacles corresponding to segments of $\partial^e P$ are free to end anywhere on the face of Σ containing the segment. We call the latter *free tentacles*, and we say that a free tentacle is a *classically allowed tentacle* if its end is in the classically allowed region $\mathcal{A}_P$. For an amoeba A , we let $\nu_{\text{AT}}(A)$ denote the number of classically allowed tentacles of A . It is clear from the above that

$$\nu_{\text{AT}}(A) \leq \#\{\text{free tentacles}\} = \text{Length}(\partial^e P)$$

and that this bound can be attained for any polytope P . Here, ‘Length’ means the length in the above sense; i.e., the diagonal face of $p\Sigma$ is scaled to have length p . As a consequence of Theorem 3 (for $m = 1$), we conclude that this maximum is asymptotically the average:

COROLLARY 4.7. *For a Delzant polytope P , we have*

$$\frac{1}{N} \mathbf{E}_{|NP} \left(\nu_{\text{AT}}(\text{Log}(Z_f)) \right) \rightarrow \text{Length}(\partial^e P).$$

Proof. Let F_1, F_2, F_3 denote the facets of $p\Sigma$ and apply Theorem 3(ii) to the 1-dimensional polytopes $P \cap \bar{F}_j$, $j = 1, 2, 3$. □

²The authors would like to thank T. Theobald for giving us permission to use this figure from his paper.

5. EXAMPLES

We now compute $b(z)$ and ψ_P for some examples of toric surfaces ($m = 2$). Recalling (10) and (135),

$$b(z) = q(z) \cdot \tau_z + p \log \left(\frac{1 + \|z\|^2}{1 + \|e^{\tau_z/2} \cdot z\|^2} \right), \quad q(z) = p \mu_\Sigma(e^{\tau_z/2} \cdot z) \in \partial P,$$

$$u_\infty = -b(z) + p \log(1 + \|z\|^2) = p \log(1 + \|e^{\tau_z/2} \cdot z\|^2) - \langle q(z), \tau_z \rangle.$$

Hence by (153),

$$\psi_P = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} u_\infty = \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \left[p \log(1 + \|e^{\tau_z/2} \cdot z\|^2) - \langle q(z), \tau_z \rangle \right]. \quad (165)$$

5.1. The square. For our first example, we let P be the unit square with vertices $\{(0, 0), (1, 0), (0, 1), (1, 1)\}$ so that $p = 2$ and $M_P = \mathbb{CP}^1 \times \mathbb{CP}^1$. Recalling that

$$\mu_\Sigma(z_1, z_2) = \left(\frac{|z_1|^2}{1 + |z_1|^2 + |z_2|^2}, \frac{|z_2|^2}{1 + |z_1|^2 + |z_2|^2} \right),$$

we see that the classically allowed region is given by

$$|z_1|^2 - 1 < |z_2|^2 < |z_1|^2 + 1,$$

as illustrated in Figure 1 in the introduction. The forbidden region consists of two subregions:

$$\mathcal{R}_F = \{(z_1, z_2) : |z_2|^2 \geq |z_1|^2 + 1\}, \quad F = \{(x_1, \frac{1}{2}) : 0 \leq x_1 \leq \frac{1}{2}\},$$

$$\mathcal{R}_{F^*} = \{(z_1, z_2) : |z_2|^2 \leq |z_1|^2 - 1\}, \quad F^* = \{(\frac{1}{2}, x_2) : 0 \leq x_2 \leq \frac{1}{2}\}.$$

Suppose that z is a point in the upper forbidden region $\mathcal{R}_F$. Write $\tau_z = (\tau_1, \tau_2)$; then $\tau_1 = 0$ since $\tau \perp T_F$. Let

$$q(z) = 2\mu_\Sigma(e^{\tau_z/2} \cdot z) = (a, 1) \in 2F \subset \partial P.$$

Writing

$$|z_1|^2 = s_1, \quad |z_2|^2 = s_2, \quad |e^{\tau_2/2} z_2|^2 = e^{\tau_2} s_2 = \tilde{s}_2,$$

we have

$$\frac{s_1}{1 + s_1 + \tilde{s}_2} = \frac{a}{2}, \quad \frac{\tilde{s}_2}{1 + s_1 + \tilde{s}_2} = \frac{1}{2}.$$

Therefore

$$s_1 = \frac{a}{1-a}, \quad \tilde{s}_2 = \frac{1}{1-a} = \frac{s_1}{a}, \quad a = \frac{s_1}{1+s_1} = \frac{|z_1|^2}{1+|z_1|^2},$$

$$e^{\tau_2} = \tilde{s}_2/s_2 = \frac{s_1}{as_2} = \frac{1+|z_1|^2}{|z_2|^2}.$$

We have

$$\log(1 + \|e^{\tau_z/2} \cdot z\|^2) = \log \left(1 + |z_1|^2 + \frac{1+|z_1|^2}{|z_2|^2} |z_2|^2 \right) = \log(1 + |z_1|^2) + \log 2,$$

$$\langle q(z), \tau_z \rangle = \left\langle \left(\frac{|z_1|^2}{1+|z_1|^2}, 1 \right), \left(0, \log \frac{1+|z_1|^2}{|z_2|^2} \right) \right\rangle = \log(1 + |z_1|^2) - \log |z_2|^2.$$

Therefore

$$u_\infty = \log |z_2|^2 + \log(1 + |z_1|^2) + \log 4.$$

We conclude that

$$\psi_P = \begin{cases} \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log(1 + |z_1|^2) & \text{for } |z_1|^2 + 1 \leq |z_2|^2 \\ 2\omega_{FS} = \frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \log(1 + |z_1|^2 + |z_2|^2) & \text{for } |z_1|^2 - 1 \leq |z_2|^2 \leq |z_1|^2 + 1 \\ \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log(1 + |z_2|^2) & \text{for } |z_2|^2 \leq |z_1|^2 - 1 \end{cases}$$

(where the third case is by symmetry). Note that ψ_P has constant rank 1 in both of the forbidden regions $\mathcal{R}_F, \mathcal{R}_{F^*}$, as indicated in Theorem 3(iii).

By the above, we also obtain

$$e^{-b(z)} = \frac{4|z_2|^2(1+|z_1|^2)}{(1+|z_1|^2+|z_2|^2)^2} \quad \text{for } |z_1|^2 + 1 \leq |z_2|^2 ,$$

and similarly for $|z_2|^2 \leq |z_1|^2 - 1$.

Remark: On the boundary $\{|z_1|^2 = |z_2|^2 - 1\}$, we have $e^{-b(z)} = 1$ as expected. On $\{|z_1| = c\}$, we have the growth rate $e^{-b(z)} \sim 1/|z_2|^2$ as $z_2 \rightarrow \infty$.

5.2. The Hirzebruch surfaces. We now consider the trapezoidal polytope of Figure 6 below.




FIGURE 6. $P = \{(0,0), (1,0), (2,0), (0,1), (1,1)\}$

Again $p = 2$, but this time $M_P \xrightarrow{\pi} \mathbb{P}^2$ is the blow up of $(0,0,1)$, i.e., M_P is the Hirzebruch surface F_1 (see [Fu]). Comparing with the case of the square, we see that the classically allowed region is given by $|z_2|^2 < |z_1|^2 + 1$, and the forbidden region coincides with the upper forbidden region $\mathcal{R}_F = \{|z_2|^2 \geq |z_1|^2 + 1\}$ from §5.1. Thus, the map $z \mapsto (\tau_z, w_z)$ is the same as before when z is in the forbidden region $\mathcal{R}_F$. Hence, $e^{-b(z)}$ and ψ_P are also the same as in §5.1 on $\mathcal{R}_F$. On the classically allowable region, $e^{-b(z)} = 1$ and $\psi_P = 2\omega_{\text{FS}}$.

5.2.1. The Hirzebruch surfaces F_n ($n \geq 2$). Let $n \geq 2$ and consider the polytope

$$P = \{(0,0), (1,0), \dots, (n+1,0), (0,1), (1,1)\} ,$$

so that $M_P = F_n$ (see [Fu]). Here $p = n+1$ and $\frac{1}{p}P$ has an interior vertex $v = (\frac{1}{n+1}, \frac{1}{n+1}) \in \Sigma^\circ$.

We see that the classically allowed region $\mu_\Sigma^{-1}(\frac{1}{p}P^\circ)$ is given by

$$|z_2|^2 < \min \left\{ \frac{|z_1|^2 + 1}{n}, \frac{1}{n-1} \right\} .$$

This time, the forbidden region consists of three subregions: $\mathcal{R}_F$, $\mathcal{R}_v$, $\mathcal{R}_{F'}$, where

$$F = \{(x_1, \frac{1}{n+1}) : 0 \leq x_1 < \frac{1}{n+1}\} , \quad F' = \{(x_1, x_2) : x_2 = \frac{1}{n}(1-x_1), \frac{1}{n+1} < x_1 \leq 1\} .$$

(See Figure 7 below.)

Suppose that z is a point in the region $\mathcal{R}_F$. Then $\tau_z = (0, \tau_2)$, as before. Let


$$q(z) = (n+1)\mu_\Sigma(e^{\tau_z/2} \cdot z) = (a, 1) \in (n+1)F \subset \partial P \quad (0 < a < 1) .$$

Again writing

$$|z_1|^2 = s_1, \quad |z_2|^2 = s_2, \quad |e^{\tau_2/2} z_2|^2 = e^{\tau_2} s_2 = \tilde{s}_2 ,$$

we have

$$\frac{s_1}{1+s_1+\tilde{s}_2} = \frac{a}{n+1} , \quad \frac{\tilde{s}_2}{1+s_1+\tilde{s}_2} = \frac{1}{n+1} .$$

FIGURE 7. $P = \{(0, 0), (1, 0), \dots, (n+1, 0), (0, 1), (1, 1)\}$

Therefore

$$s_1 = \frac{a}{n-a}, \quad \tilde{s}_2 = \frac{1}{n-a} = \frac{s_1}{a}, \quad a = \frac{ns_1}{1+s_1} = \frac{n|z_1|^2}{1+|z_1|^2},$$

$$e^{\tau_2} = \tilde{s}_2/s_2 = \frac{s_1}{as_2} = \frac{1+|z_1|^2}{n|z_2|^2}.$$

In particular,

$$a < 1 \Leftrightarrow |z_1|^2 < \frac{1}{n-1}$$

and therefore

$$\mathcal{R}_F = \left\{ (z_1, z_2) : |z_2|^2 \geq \frac{|z_1|^2 + 1}{n}, |z_1|^2 < \frac{1}{n-1} \right\}.$$

We have

$$\log(1 + \|e^{\tau_z/2} \cdot z\|^2) = \log \left(1 + |z_1|^2 + \frac{1+|z_1|^2}{n|z_2|^2} |z_2|^2 \right) = \log(1 + |z_1|^2) + \log \frac{n+1}{n},$$

$$\langle q(z), \tau_z \rangle = \left\langle \left(\frac{n|z_1|^2}{1+|z_1|^2}, 1 \right), \left(0, \log \frac{1+|z_1|^2}{n|z_2|^2} \right) \right\rangle = \log(1 + |z_1|^2) - \log |z_2|^2 - \log n.$$

Therefore

$$u_\infty = \log |z_2|^2 + n \log(1 + |z_1|^2) + (n+1) \log(n+1) - n \log n.$$

Hence,

$$\left. \begin{aligned} \psi_P &= n \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log(1 + |z_1|^2) \\ e^{-b(z)} &= \frac{(n+1)^{n+1}}{n^n} \frac{|z_2|^2 (1 + |z_1|^2)^n}{(1 + |z_1|^2 + |z_2|^2)^{n+1}} \end{aligned} \right\} \quad \text{for } z \in \mathcal{R}_F.$$

Now suppose that z is a point in $\mathcal{R}_{F'}$. Since $\tau_z \perp T_{F'}$, we can write $\tau_z = (\tau_1, n\tau_1)$. Let

$$\frac{1}{n+1} q(z) = \mu_\Sigma(e^{\tau_z/2} \cdot z) = \left(x_1, \frac{1}{n}(1 - x_1) \right) \in F'. \quad (166)$$

As before, we write

$$s_1 = |z_1|^2, \quad s_2 = |z_2|^2, \quad \tilde{s}_1 = |e^{\tau_1/2} z_1|^2 = e^{\tau_1} s_1, \quad \tilde{s}_2 = |e^{n\tau_1/2} z_2|^2 = e^{n\tau_1} s_2.$$

By (166), we have

$$\frac{\tilde{s}_1}{1 + \tilde{s}_1 + \tilde{s}_2} = x_1, \quad \frac{\tilde{s}_2}{1 + \tilde{s}_1 + \tilde{s}_2} = \frac{1}{n}(1 - x_1).$$

Solving for $\tilde{s}_1, \tilde{s}_2$, we obtain

$$\tilde{s}_1 = \frac{n}{n-1} \frac{x_1}{1-x_1}, \quad \tilde{s}_2 = \frac{1}{n-1}. \quad (167)$$

Therefore, $e^{n\tau_1} = \tilde{s}_2/s_2 = |z_2|^{-2}/(n-1)$, so we have

$$\tau_1 = -\frac{1}{n} \log(n-1)|z_2|^2, \quad \tilde{s}_1 = \frac{|z_1|^2}{(n-1)^{1/n}|z_2|^{2/n}}.$$

Thus,

$$\log(1 + \|e^{\tau_z/2} \cdot z\|^2) = \log(1 + \tilde{s}_1 + \tilde{s}_2) = \log\left(\frac{n}{n-1} + \frac{|z_1|^2}{(n-1)^{1/n}|z_2|^{2/n}}\right).$$

By (166),

$$\langle q(z), \tau_z \rangle = (n+1) \left\langle \left(x_1, \frac{1}{n}(1-x_1)\right), (\tau_1, n\tau_1) \right\rangle = (n+1)\tau_1 = -\frac{n+1}{n} \log(n-1)|z_2|^2.$$

Hence by (165),

$$\psi_P = (n+1) \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log\left(\frac{n}{n-1} + \frac{|z_1|^2}{(n-1)^{1/n}|z_2|^{2/n}}\right) \quad \text{for } z \in \mathcal{R}_{F'}.$$

(Note that ψ_P has constant rank 1 on $\mathcal{R}_{F'}$ as indicated by Theorem 3(iii), since $\psi_P = (n+1)g^*\omega_{\mathbb{CP}^1}$ on $\mathcal{R}_{F'}$, where g is the multi-valued holomorphic map to $\mathbb{CP}^1$ given by $g(z_1, z_2) = (c_n z_1, z_2^{1/n})$.)

By Theorem 3(iii), we know that $\psi_P = 0$ on $\mathcal{R}_v$. To complete the description of ψ_P , it remains to describe the regions $\mathcal{R}_{F'}$ and $\mathcal{R}_v$. We note that a forbidden point z lies in $\mathcal{R}_{F'}$ if and only if $x_1 > \frac{1}{n+1}$. By (167), this is equivalent to $\tilde{s}_1 > \frac{1}{n-1}$, or

$$|z_2|^2 < (n-1)^{n-1}|z_1|^{2n}.$$

Therefore

$$\mathcal{R}_{F'} = \left\{ (z_1, z_2) : \frac{1}{n-1} \leq |z_2|^2 < (n-1)^{n-1}|z_1|^{2n}, \quad |z_1|^2 > \frac{1}{n-1} \right\}.$$

This leaves us with

$$\mathcal{R}_v = \left\{ (z_1, z_2) : |z_2|^2 \geq (n-1)^{n-1}|z_1|^{2n}, \quad |z_1|^2 \geq \frac{1}{n-1} \right\}.$$

To summarize:

$$\psi_P = \begin{cases} (n+1) \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log(1 + |z_1|^2 + |z_2|^2) & \text{for } |z_2|^2 < \min\left\{\frac{|z_1|^2+1}{n}, \frac{1}{n-1}\right\} \\ n \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log(1 + |z_1|^2) & \text{for } |z_2|^2 \geq \frac{|z_1|^2+1}{n}, \quad |z_1|^2 < \frac{1}{n-1} \\ (n+1) \frac{\sqrt{-1}}{2\pi} \partial \bar{\partial} \log\left(\frac{n}{n-1} + \frac{|z_1|^2}{(n-1)^{1/n}|z_2|^{2/n}}\right) & \text{for } \frac{1}{n-1} \leq |z_2|^2 < (n-1)^{n-1}|z_1|^{2n} \\ 0 & \text{for } |z_2|^2 \geq (n-1)^{n-1}|z_1|^{2n}, \quad |z_1|^2 \geq \frac{1}{n-1} \end{cases}$$

5.3. Remarks on the behavior at transition points. So far, we have studied the uniform asymptotics of $\Pi_{|NP}(z, z)$ on compact subsets of the classically allowed region and on compact sets without transition points in the forbidden region. We now address a few words to the asymptotics at the boundary of the classically allowed region, which is its caustic set. For background on oscillatory integrals and caustics, we refer to [AGV].

We thus consider $\Pi_{|NP}(z, z)$ as an oscillatory integral (denoted $K_N(z)$ in (72)) on $(\mathbb{C}^*)^m$. Since it is $\mathbf{T}^m$ invariant we may regard it more simply as an oscillatory integral in $z = e^\rho$ on $\mathbb{R}_+^m$ or on $p\Sigma$, with the phase

Ψ of (73) regarded as a function on $\mathbf{T}^m \times M_P \times \mathbb{R}_+^m$. The asymptotics of $\Pi_{|NP}(z, z)$ are determined by the component

$$C^0 = \{(0, w; e^\rho) \in \mathbb{R}_+^m \times M_P : \mu_{p\Sigma}(e^\rho) = \mu_P(w)\}, \quad (168)$$

of the critical set on which $\varphi = 0$, so we may (and will) ignore other critical points which may occur. The map $(0, w; e^\rho) \rightarrow w$ gives a natural identification of C^0 with M_P .

As is standard in the theory of oscillatory integrals depending on parameters, we define the map

$$\iota_\Psi : C^0 \rightarrow T^*\mathbb{R}_+^m, \quad \iota_\Psi(0, w; e^\rho) = (e^\rho, d_\rho \Psi), \quad (169)$$

and we denote the image by $\Lambda^0 = \iota_\Psi(C^0)$. Over the set of parameters z for which the critical point set C_z^0 is a non-degenerate critical manifold, ι_Ψ is a fibration and Λ^0 is a Lagrangean manifold. The set of $z = e^\rho$ for which the phase is degenerate is known as the caustic set. Under the identification of $C^0 \sim M_P$, we have $\frac{1}{2}d_z \Psi(w; e^\rho) \equiv 0$ on C^0 , hence Λ^0 is the zero section of T^*P . Identifying ι_Ψ with the map $\mu_P \times 0$, we find that the caustic set is the boundary of the classically allowed region.

Uniform asymptotic expansions of $\Pi_{|NP}(z, z)$ in the vicinity of the caustic could be obtained with a careful study of the singularities of the moment map μ_P . The uniform expansion would give a smooth transition in a boundary layer from the N^m mass density in the interior of P (the illuminated or allowed region) to the $e^{-Nb(z)}$ mass density in the exterior (the shadow or forbidden region). The order of the asymptotics along the caustic depends on two competing effects: on the one hand, oscillatory integrals whose (analytic) phase has a degenerate critical point decay more slowly than in the non-degenerate case (by the index of the singularity at the critical point, see [AGV]). On the other hand, the principal term of the amplitude may vanish on the caustic, causing more rapid decay. We will be content to illustrate this in the simplest example.

5.3.1. *Example:* $P = [0, 1], p = 2$. We put $m = 1, P = [0, 1], p = 2, p\Sigma = [0, 2]$. Thus, $M_P = \mathbb{CP}^1$. The only interior facet is $F = \{1\}$; recalling (73), our phase is:

$$\Psi(w, \varphi; z) = \log \frac{1 + e^{-i\varphi}|w|^2}{1 + |w|^2} + 2 \log \frac{1 + e^{i\varphi}|z|^2}{1 + |z|^2}. \quad (170)$$

We are interested in the asymptotics of $\Pi_{|NP}(z, z)$ when

$$\mu_{2\Sigma}(z) = \frac{2|z|^2}{1 + |z|^2} = 1 \iff |z| = 1.$$

We claim that $\Pi_{|NP}(z, z) \sim CN^1$ when $|z| = 1$.

Since $\mu_P(w) = \frac{|w|^2}{1 + 2|w|^2}$, we have $\mu_P(w) = \{1\} \iff w = \infty$. To determine the asymptotics of $\Pi_{|NP}(z, z)$ at $|z| = 1$, we put $\eta = \frac{1}{w} \in \mathbb{C}$ and consider the Taylor expansion of the phase at $\eta = 0, \varphi = 0$. We change to polar coordinates $\eta = re^{i\theta}$ around the fixed point of the toric S^1 action. The phase and amplitude are independent of θ and the principal part of the Taylor series (at $|z| = 1$) has the form:

$$\Psi(w, \varphi; 1) \sim i\varphi r^2 - \frac{1}{4}\varphi^2 + O(|\varphi|^3 + |\varphi|^2|r|^2 + |\varphi||r|^4).$$

By a general result [AGV], the asymptotics of (generic) oscillatory integrals are determined by the principal part of the Taylor expansion of the phase at the critical point, which gives the model integral

$$I_N = \int_{-\pi}^{\pi} \int_0^{\infty} e^{N(i\varphi r^2 - \frac{1}{4}\varphi^2)} A(r, \varphi) r dr d\varphi,$$

where A is an amplitude decaying sufficiently rapidly at infinity. Here, we also used that the amplitude of $\Pi_{|NP}(z, z)$ is simply the volume density in our local coordinates (φ, r, θ) . We observe that the phase $f(r, \varphi) = i\varphi r^2 - \frac{1}{4}\varphi^2$ is *weighted homogeneous* in the sense that

$$f(N^{1/4}r, N^{1/2}\varphi) = Nf(r, \varphi).$$

We also observe that the amplitude vanishes to order one as $r \rightarrow 0$. We change variables as indicated to obtain that

$$\Pi_{|NP}(1, 1) \sim N^2 I_N = N^2 \int_{\mathbb{R}} \int_{\mathbb{R}} N^{-1} e^{(i\varphi r^2 + C(z)\varphi^2)} \rho(N^{-1/4}r, N^{-1/2}\varphi) r dr d\varphi \sim CN,$$

as claimed.

Alternatively, we observe that the index of oscillation of the phase with $|z| = 1$ equals $-3/4$. This agrees with the general result (under assumptions satisfied here) that the index of oscillation equals the ‘remoteness’ of the Newton polygon of the phase ([AGV], §II.6, Theorem 4), i.e. the reciprocal of the value of t such that the ray (t, t) , $t > 0$ intersects the boundary of the Newton polygon. The index of the singularity is therefore $\delta = 1/4$, and therefore $I_N \sim CN^{-1}$.

REFERENCES

- [Ag] S. Agmon, *Lectures on exponential decay of solutions of second-order elliptic equations: bounds on eigenfunctions of N -body Schrödinger operators*. Mathematical Notes 29, Princeton University Press, Princeton, NJ, 1982.
- [AGV] V. I. Arnold, S. M. Gusein-Zade and A. N. Varchenko, *Singularities of differentiable maps. Vol. II. Monodromy and asymptotics of integrals*, Monographs in Math. 83, Birkhäuser, Boston, 1988.
- [At] M.F. Atiyah, Angular momentum, convex polyhedra and algebraic geometry, *Proc. Edinburgh Math. Soc.* 26 (1983), 121–138.
- [Au] M. Audin, *The Topology of Torus Actions on Symplectic Manifolds*, Progress in Math. 93, Birkhäuser, Boston, 1991.
- [BT] E. Bedford and B. A. Taylor, A new capacity for plurisubharmonic functions, *Acta Math.* 149 (1982), 1–40.
- [Be] D. N. Bernstein, The number of roots of a system of equations, *Functional Anal. Appl.* 9 (1975), 183–185.
- [BSZ] P. Bleher, B. Shiffman and S. Zelditch, Universality and scaling of correlations between zeros on complex manifolds, *Invent. Math.* 142 (2000), 351–395.
- [BG] L. Boutet de Monvel and V. Guillemin, *The spectral theory of Toeplitz operators*. Annals of Mathematics Studies, 99. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1981.
- [BS] L. Boutet de Monvel and J. Sjöstrand, Sur la singularité des noyaux de Bergman et de Szegő, *Asterisque* 34–35 (1976), 123–164.
- [BV] M. Brion and M. Vergne, Lattice points in simple polytopes, *J. Amer. Math. Soc.* 10 (1997), 371–392.
- [Da] V. Danilov, The geometry of toric varieties, *Russian Math. Surveys* 33 (1978), 97–154.
- [De] T. Delzant, Hamiltoniens périodiques et image convexe de l’application moment, *Bull. Soc. Math. France* 116 (1988), 315–339.
- [Fan] X. Fan, The necessary and sufficient conditions for Lipschitz local homeomorphism, *Chinese Ann. Math. Ser. B* 13 (1992), 40–45.
- [Fe] H. Federer, *Geometric measure theory*, Springer-Verlag, New York, 1969.
- [FPT] M. Forsberg, M. Passare and A. Tsikh, Laurent determinants and arrangements of hyperplane amoebas, *Adv. in Math.* 151 (2000), 45–70.
- [Fu] W. Fulton, *Introduction to Toric Varieties*, Ann. Math. Studies 131, Princeton Univ. Press, Princeton (1993).
- [GKZ] I. M. Gelfand, M. M. Kapranov and A. V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Mathematics: Theory and Applications, Birkhäuser, Boston, 1994.
- [Gu1] V. Guillemin, *Moment Maps and Combinatorial Invariants of Hamiltonian T^n -Spaces*, Progress in Math. 122, Birkhäuser, Boston, 1994.
- [Gu2] V. Guillemin, Star products on compact pre-quantizable symplectic manifolds, *Lett. Math. Phys.* 35 (1995), 85–89.
- [GS] V. Guillemin and S. Sternberg, Geometric quantization and multiplicities of group representations, *Invent. Math.* 67 (1982), 515–538.
- [Hö] L. Hörmander, *The Analysis of Linear Partial Differential Operators, I*, Springer Verlag, N.Y., 1983.
- [HS] B. Huber and B. A. Sturmfels, polyhedral method for solving sparse polynomial systems, *Math. Comp.* 64 (1995), 1541–1555.
- [KP] A. G. Khovanskii and A.V. Pukhlikov, A Riemann-Roch theorem for integrals and sums of quasi-polynomials over virtual polytopes, *St. Petersburg Math. J.* 6 (1993), 789–812.
- [Kl] M. Klimek, *Pluripotential Theory*, London Math. Soc. Monographs, New Series 6, Oxford University Press, New York, 1991.
- [Ko1] A. G. Kouchnirenko, Polyèdres de Newton et nombres de Milnor, *Invent. Math.* 32 (1976), 1–31.
- [Ko2] A. G. Kouchnirenko, Newton Polytopes and the Bezout theorem, *Functional Anal. Appl.* 10 (1976), 233–235.
- [MR] G. Malajovich and J. M. Rojas, Random Sparse Polynomial Systems, e-print archive, math.NA/0012104.
- [MS] A. Melin and J. Sjöstrand, Fourier integral operators with complex-valued phase functions. *Fourier integral operators and partial differential equations* (Colloq. Internat., Univ. Nice, Nice, 1974), Lecture Notes in Math., Vol. 459, Springer, Berlin, 1975, pp. 120–223.
- [Mi1] G. Mikhalkin, Real algebraic curves, the moment map and amoebas, *Ann. Math.* 151 (2000), 309–326.
- [Mi2] G. Mikhalkin, Amoebas of algebraic varieties, e-print archive, math.AG/0108225.

- [Oda] T. Oda, *Convex Bodies and Algebraic Geometry*, Ergebnisse 15, Springer-Verlag, New York (1988).
- [PR] M. Passare and H. Rullgård, Ameobas, Monge-Ampere measures and triangulations of the Newton polytope (preprint, Stockholm University, 2001).
- [RT] Reshetikhin, N.; Takhtajan, L. A. Deformation quantization of Kähler manifolds. *L. D. Faddeev's Seminar on Mathematical Physics*, 257–276, Amer. Math. Soc. Transl. Ser. 2, 201, Amer. Math. Soc., Providence, RI, 2000.
- [Ro] J. M. Rojas, On the average number of real roots of certain random sparse polynomial systems. In *The mathematics of numerical analysis (Park City, UT, 1995)*, 689–699, Lectures in Appl. Math. 32, Amer. Math. Soc., Providence, RI, 1996.
- [SZ1] B. Shiffman and S. Zelditch, Distribution of zeros of random and quantum chaotic sections of positive line bundles, *Comm. Math. Phys.* 200 (1999), 661–683.
- [SZ2] B. Shiffman and S. Zelditch, Random polynomials with prescribed Newton polytope, II: non-Delzant polytopes (in preparation).
- [St] B. Sturmfels, On the number of real roots of a sparse polynomial system, *Hamiltonian and gradient flows, algorithms and control*, Fields Inst. Commun. 3, Amer. Math. Soc., Providence, RI, 1994, pp. 137–143.
- [Th] T. Theobald, Computing amoebas (preprint, Technische Universität München, Report TUM-M0200, 2002).
- [Ve] J. Verschelde, Toric Newton method for polynomial homotopies, *J. Symbolic Computation* 29 (2000), 777–793.
- [Ze] S. Zelditch, Szegő kernels and a theorem of Tian, *IMRN* 6 (1998), 317–331.

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD 21218, USA

E-mail address: `shiffman@math.jhu.edu`, `zelditch@math.jhu.edu`